

Anti-Resistin/RETN Antibody Picoband™

Catalog Number: PB9412

About RETN

Resistin, also known as adipose tissue-specific secretory factor (ADSF) or C/EBP-epsilon-regulated myeloid-specific secreted cysteine-rich protein (XCP1) is a cysteine-rich protein that in humans is encoded by the RETN gene. The resistin gene comprises 4 exons, the first of which is untranslated, and spans approximately 1,750 bp. The human resistin gene is localized to a cloned fragment of human chromosome 19. In primates, pigs and dogs, resistin is secreted by immune and epithelial cells while in rodents, it is secreted by adipose tissue. Resistin is a cytokine whose physiologic role has been the subject of much controversy regarding its involvement with obesity and type II diabetes mellitus.

Overview

| | |
|----------------------|---|
| Product Name | Anti-Resistin/RETN Antibody Picoband™ |
| Reactive Species | Human |
| Description | Boster Bio Anti-Resistin/RETN Antibody Picoband™ catalog # PB9412. Tested in ELISA, WB applications. This antibody reacts with Human. |
| Application | ELISA, WB |
| Clonality | Polyclonal |
| Formulation | Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3. |
| Storage Instructions | Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles. |
| Host | Rabbit |
| Uniprot ID | Q9HD89 |

Technical Details

| | |
|-------------------------------|---|
| Immunogen | E. coli-derived human Resistin recombinant protein (Position: K19-P108). Human Resistin shares 58.6% amino acid (aa) sequence identity with mouse Resistin. |
| Predicted Reactive Species | Bovine |
| Recommended Detection Systems | Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot. |
| Cross Reactivity | No cross-reactivity with other proteins. |
| Isotype | Rabbit IgG |
| Form | Lyophilized |
| Concentration | Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. |

| | |
|---------------------|---|
| Purification | Immunogen affinity purified. |
| Suggested Dilutions | <p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.5ug/ml, Human , -</p> <p>ELISA , 0.1-0.5ug/ml, Human</p> |

Anti-Resistin/RETN Antibody Picoband™ (PB9412) Images



Figure 1. Western blot analysis of Resistin using anti-Resistin antibody (PB9412).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours.

Lane 1: Recombinant Human Resistin Protein 0.5ng.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Resistin antigen affinity purified polyclonal antibody (Catalog # PB9412) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Resistin at approximately 11 kDa. The expected band size for Resistin is at 11 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Resistin/RETN Antibody™