

Anti-RAB10 Antibody Picoband™

Catalog Number: PB9788

About RAB10

Ras-related protein Rab-10 is a protein that in humans is encoded by the RAB10 gene. RAB10 belongs to the RAS superfamily of small GTPases. And RAB proteins localize to exocytic and endocytic compartments and regulate intracellular vesicle trafficking. The RAB10 gene is mapped to chromosome 2p23.1-p22.3 by radiation hybrid analysis. It was found that the purified recombinant GAP domain of human AS160 showed GAP activity with RAB2A, RAB8A, RAB10, and RAB14, but not with 14 other RABs. Immunoblot analysis showed that these RABs associated with Glut4-positive vesicles in mouse adipocytes. Thereby, it concluded that AK160 functions as a RAB GAP and that RABs may participate in GLUT4 translocation.

Overview

Product Name	Anti-RAB10 Antibody Picoband™
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-RAB10 Antibody Picoband™ catalog # PB9788. Tested in WB applications. This antibody reacts with Human, Mouse, Rat.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P61026

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human RAB10, identical to the related mouse sequence, and different from the related rat sequence by one amino acid.
Predicted Reactive Species	Hamster
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat

Anti-RAB10 Antibody Picoband™ (PB9788) Images




Figure 1. Western blot analysis of RAB10 using anti-RAB10 antibody (PB9788).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human MCF-7 whole cell lysates,

Lane 2: human Hela whole cell lysates,

Lane 3: human A549 whole cell lysates,

Lane 4: rat brain tissue lysates,

Lane 5: rat C6 whole cell lysates,

Lane 6: mouse brain tissue lysates,

Lane 7: mouse NIH/3T3 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes.

Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-RAB10 antigen affinity purified polyclonal antibody (Catalog # PB9788) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for RAB10 at approximately 23 kDa. The expected band size for RAB10 is at 23 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-RAB10 Antibody™