

Anti-Rab18 Antibody Picoband™

Catalog Number: PB9791

About RAB18

Ras-related protein Rab-18 is a protein that in humans is encoded by the RAB18 gene. It is a ubiquitously expressed protein with particularly high expression in the brain. The protein encoded by this gene is a member of a family of Ras-related small GTPases that regulate membrane trafficking in organelles and transport vesicles. Knockdown studies in zebrafish suggest that this protein may have a role in eye and brain development. Mutations in this gene are associated with Warburg micro syndrome type 3. Alternatively spliced transcript variants have been found for this gene.

Overview

Product Name	Anti-Rab18 Antibody Picoband™
Reactive Species	Human, Rat
Description	Boster Bio Anti-Rab18 Antibody Picoband™ catalog # PB9791. Tested in WB applications. This antibody reacts with Human, Rat.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q9NP72

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human Rab18, identical to the related mouse sequence, and different from the related rat sequence by one amino acid.
Predicted Reactive Species	Bovine
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.1-0.5ug/ml, Human, Rat

Anti-Rab18 Antibody Picoband™ (PB9791) Images



Figure 1. Western blot analysis of Rab18 using anti-Rab18 antibody (PB9791). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. Lane 1: Rat Testis Tissue Lysate at 50ug, Lane 2: Rat Lung Tissue Lysate at 50ug, Lane 3: 293T Whole Cell Lysate at 40ug, Lane 4: HELA Whole Cell Lysate at 40ug, Lane 5: HEPG2 Whole Cell Lysate at 40ug. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Rab18 antigen affinity purified polyclonal antibody (Catalog # PB9791) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Rab18 at approximately 23 kDa. The expected band size for Rab18 is at 23 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Rab18 Antibody™