

Anti-TGF beta Receptor I/TGFB1 Antibody Picoband™

Catalog Number: PB9828

About TGFB1

Transforming growth factor, beta receptor I is a TGF beta receptor. TGFB1 is its human gene. The protein encoded by this gene forms a heteromeric complex with type II TGF-beta receptors when bound to TGF-beta, transducing the TGF-beta signal from the cell surface to the cytoplasm. Mutations in this gene have been associated with Loeys-Dietz aortic aneurysm syndrome (LDAS). TGFB1 regulates cell cycle progression by a unique signaling mechanism that involves its binding to TGFB2 and activation of TGFB1. Both are transmembrane serine/threonine receptor kinases. The TGFB1 receptor may be inactivated in many of the cases of human tumor cells refractory to TGF-beta-mediated cell cycle arrest. Vellucci and Reiss (1997) reported that the TGFB1 gene is approximately 31 kb long and contains 9 exons. The organization of the segment of the gene that encodes the C-terminal portion of the serine/threonine kinase domain appears to be highly conserved among members of the gene family.

Overview

Product Name	Anti-TGF beta Receptor I/TGFB1 Antibody Picoband™
Reactive Species	Human
Description	Boster Bio Anti-TGF beta Receptor I/TGFB1 Antibody Picoband™ catalog # PB9828. Tested in WB applications. This antibody reacts with Human.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P36897

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human TGFB1, identical to the related mouse and rat sequences.
Predicted Reactive Species	Bovine, Chicken, Horse, Monkey, Rabbit
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG

Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.1-0.5ug/ml, Human

Anti-TGF beta Receptor I/TGFBR1 Antibody Picoband™ (PB9828) Images




Figure 1. Western blot analysis of TGFBR1/Tgf Beta Receptor I using anti-TGFBR1/Tgf Beta Receptor I antibody (PB9828).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysates,

Lane 2: human SW579 whole cell lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TGFBR1/Tgf Beta Receptor I antigen affinity purified polyclonal antibody (Catalog # PB9828) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for TGFBR1/Tgf Beta Receptor I at approximately 50-56KD. The expected band size for TGFBR1/Tgf Beta Receptor I is at 56KD.

3 Publications Citing This Product

1. PubMed ID: 11780431, Effects of AT1 receptor antagonist, losartan, on rat hepatic fibrosis induced by CCI4
2. PubMed ID: 26279755, Different localization and expression of protein kinase C-beta in kidney cortex of diabetic nephropathy mice and its role in telmisartan treatment
3. PubMed ID: 21464546, Effects of interferon-alpha on expression of hepatic stellate cell and transforming growth factor- β 1 and β -smooth muscle actin in rats with hepatic fibrosis

Visit bosterbio.com/anti-tgfr1-picoband-trade-antibody-pb9828-boster.html to see all 3 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-TGF beta Receptor I/TGFBR1 Antibody [™]