

Anti-TGF beta Receptor II/TGFR2 Antibody Picoband™

Catalog Number: PB9829

About TGFR2

TGFR2 (transforming growth factor, beta receptor II (70/80kDa)), also known as TGF-beta receptor type-2, TGFR-2, TGF-beta type II receptor, Transforming growth factor-beta receptor type II (TGF-beta receptor type II, TbetaR-II), is a member of the Ser/Thr protein kinase family and the TGFB receptor subfamily. A TGFR2 cDNA encodes a deduced 565-amino acid protein with a calculated molecular mass of approximately 60 kD in length. The encoded protein is a transmembrane protein that has a protein kinase domain, forms a heterodimeric complex with another receptor protein, and binds TGF-beta. This receptor/ligand complex phosphorylates proteins, which then enter the nucleus and regulate the transcription of a subset of genes related to cell proliferation. Mutations in this gene have been associated with Marfan syndrome, Loeys-Deitz aortic aneurysm syndrome, Osler-Weber-Rendu syndrome, and the development of various types of tumors. Alternatively spliced transcript variants encoding different informs have been characterized.

Overview

Product Name	Anti-TGF beta Receptor II/TGFR2 Antibody Picoband™
Reactive Species	Human
Description	Boster Bio Anti-TGF beta Receptor II/TGFR2 Antibody Picoband™ catalog # PB9829. Tested in WB applications. This antibody reacts with Human.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P37173

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human TGFR2, different from the related mouse sequence by five amino acids, and from the related rat sequence by eight amino acids.
Predicted Reactive Species	Bovine, Chicken, Horse, Monkey, Rabbit
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins

Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.1-0.5ug/ml, Human

Anti-TGF beta Receptor II/TGFR2 Antibody Picoband™ (PB9829) Images




Figure 1. Western blot analysis of TGFR2 using anti-TGFR2 antibody (PB9829).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 40 ug of sample under reducing conditions.

Lane 1: MCF-7 Whole Cell Lysate,
Lane 2: SW620 Whole Cell Lysate,
Lane 3: 22RV1 Whole Cell Lysate,
Lane 4: HELA Whole Cell Lysate.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TGFR2 antigen affinity purified polyclonal antibody (Catalog # PB9829) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for TGFR2 at approximately 85 kDa. The expected band size for TGFR2 is at 85 kDa.

2 Publications Citing This Product

1. PubMed ID: 26839890, Mechanism Investigation of the Improvement of Chang Run Tong on the Colonic Remodeling in Streptozotocin-Induced Diabetic Rats
2. PubMed ID: 27019660, Effect of Kuijie Granule on the Expression of TGF- β /Smads Signaling Pathway in Patients with Ulcerative Colitis

Visit bosterbio.com/anti-tgfr2-picoband-trade-antibody-pb9829-boster.html to see all 2 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-TGF beta Receptor II/TGFR2 Antibody™