

Anti-TIMP3 Antibody Picoband™

Catalog Number: PB9831

About TIMP3

Metalloproteinase inhibitor 3 is a protein that in humans is encoded by the TIMP3 gene. It is mapped to 22q12.1-q13.2. This gene belongs to the tissue inhibitor of metalloproteinases gene family. The proteins encoded by this gene family are inhibitors of the matrix metalloproteinases, a group of peptidases involved in degradation of the extracellular matrix (ECM). Expression of this gene is induced in response to mitogenic stimulation and this netrin domain-containing protein is localized to the ECM. Mutations in this gene have been associated with the autosomal dominant disorder Sorsby's fundus dystrophy.

Overview

Product Name	Anti-TIMP3 Antibody Picoband™
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti- TIMP3 Antibody Picoband™ catalog # PB9831. Tested in ELISA, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P35625

Technical Details

Immunogen	E. coli-derived human TIMP3 recombinant protein (Position: C24-P211). Human TIMP3 shares 97.9% and 97.3% amino acid (aa) sequence identity with mouse and rat TIMP3, respectively.
Predicted Reactive Species	Bovine, Chicken, Horse, Monkey, Rabbit
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: ELISA , 0.1-0.5ug/ml, Human, - Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat

Anti-TIMP3 Antibody Picoband™ (PB9831) Images




Figure 1. Western blot analysis of TIMP3 using anti-TIMP3 antibody (PB9831).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours.

Lane 1: Rat Kidney Tissue Lysate at 50ug,

Lane 2: Mouse Ovary Tissue Lysate at 50ug,

Lane 3: HELA Whole Cell Lysate at 40ug,

Lane 4: MCF-7 Whole Cell Lysate at 40ug,

Lane 5: SMMC Whole Cell Lysate at 40ug.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes.

Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT.

The membrane was incubated with rabbit anti-TIMP3 antigen affinity purified polyclonal antibody (Catalog # PB9831) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed

with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit

(Catalog # EK1002) with Tanon 5200 system. A specific band was detected for TIMP3 at approximately 34 kDa. The expected band size for TIMP3 is at 34 kDa.

3 Publications Citing This Product

1. PubMed ID: 33649809, Nie L,Liu M,Chen J,Wu Q,Li Y,Yi J,Zheng X,Zhang J,Chu C,Yang J.Hydrogen sulfide ameliorates doxorubicin-induced myocardial fibrosis in rats via the PI3K/AKT/mTOR pathway.Mol Med Rep.2021 Apr;23(4):299.doi:10.3892/mmr.2021.11938.Epub 2021 Mar 2.PMID:336498

2. PubMed ID: 23819566, Promoter methylation and expression of TIMP3 gene in gastric cancer

3. PubMed ID: 17657847, Bai Yx, Yi JI, Li Jf, Sui H. World J Gastroenterol. 2007 Jul 28;13(28):3883-5. Clinicopathologic Significance Of Bag1 And Timp3 Expression In Colon Carcinoma.

Visit bosterbio.com/anti-timp3-picoband-trade-antibody-pb9831-boster.html to see all 3 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-TIMP3 Antibody™