

Anti-Nucleobindin 2/NUCB2 Antibody Picoband™

Catalog Number: PB9876

About Nucb2

Nucleobindin-2 is a calcium-binding EF-hand protein that in humans is encoded by the NUCB2 gene. By exon trapping and sequence analysis, the NUCB2 sequence has been identified on chromosome 11p15-p14. This gene encodes a protein with a suggested role in calcium level maintenance, eating regulation in the hypothalamus, and release of tumor necrosis factor from vascular endothelial cells.

Overview

Product Name	Anti-Nucleobindin 2/NUCB2 Antibody Picoband™
Reactive Species	Mouse, Rat
Description	Boster Bio Anti-Nucleobindin 2/NUCB2 Antibody Picoband™ catalog # PB9876. Tested in ELISA, IHC, WB applications. This antibody reacts with Mouse, Rat.
Application	ELISA, IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P81117

Technical Details

Immunogen	E. coli-derived mouse Nucleobindin 2 recombinant protein (Position: V25-L106). Mouse Nucleobindin 2 shares 85.4% and 97.6% amino acid (aa) sequence identity with human and rat Nucleobindin 2, respectively.
Predicted Reactive Species	Bovine
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Mouse, Rat, By Heat ELISA , 0.1-0.5ug/ml, Mouse, - Western blot, 0.1-0.5ug/ml, Mouse

Anti-Nucleobindin 2/NUCB2 Antibody Picoband™ (PB9876) Images



Figure 1. Western blot analysis of Nucleobindin 2 using anti-Nucleobindin 2 antibody (PB9876).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: HEPA whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Nucleobindin 2 antigen affinity purified polyclonal antibody (Catalog # PB9876) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Nucleobindin 2 at approximately 55 kDa. The expected band size for Nucleobindin 2 is at 50 kDa.




Figure 2. IHC analysis of Nucleobindin 2 using anti-Nucleobindin 2 antibody (PB9876).

Nucleobindin 2 was detected in a paraffin-embedded section of mouse gaster tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 ug/ml rabbit anti-Nucleobindin 2 Antibody (PB9876) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.




Figure 3. IHC analysis of Nucleobindin 2 using anti-Nucleobindin 2 antibody (PB9876).

Nucleobindin 2 was detected in a paraffin-embedded section of rat gaster tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 ug/ml rabbit anti-Nucleobindin 2 Antibody (PB9876) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your



reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Nucleobindin 2/NUCB2 Antibody [™]