

Anti-Prokineticin 1/PROK1 Antibody Picoband™

Catalog Number: PB9881

About PROK1

Prokineticin-1 is also known as EG-VEGF. The International Radiation Hybrid Mapping Consortium mapped the PROK1 gene to chromosome 1. The protein encoded by this gene induces proliferation, migration, and fenestration (the formation of membrane discontinuities) in capillary endothelial cells derived from endocrine glands. It has little or no effect on a variety of other endothelial and non-endothelial cell types. Its expression is restricted to the steroidogenic glands (ovary, testis, adrenal, and placenta), is induced by hypoxia, and often complementary to the expression of vascular endothelial growth factor (VEGF), suggesting that these molecules function in a coordinated manner.

Overview

Product Name	Anti-Prokineticin 1/PROK1 Antibody Picoband™
Reactive Species	Human
Description	Boster Bio Anti-Prokineticin 1/PROK1 Antibody Picoband™ catalog # PB9881. Tested in ELISA, WB applications. This antibody reacts with Human.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P58294

Technical Details

Immunogen	E. coli-derived human Prokineticin 1 recombinant protein (Position: A20-F105). Human Prokineticin 1 shares 88.4% and 91.9% amino acid (aa) sequence identity with mouse and rat Prokineticin 1, respectively.
Predicted Reactive Species	Chicken
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: ELISA , 0.1-0.5ug/ml, Human, - Western blot, 0.1-0.5ug/ml, Human

Anti-Prokineticin 1/PROK1 Antibody Picoband™ (PB9881) Images




Figure 1. Western blot analysis of Prokineticin 1 using anti-Prokineticin 1 antibody (PB9881). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: MCF-7 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Prokineticin 1 antigen affinity purified polyclonal antibody (Catalog # PB9881) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Prokineticin 1 at approximately 20 kDa. The expected band size for Prokineticin 1 is at 12 kDa.

1 Publications Citing This Product

1. PubMed ID: 21139695, Administration of pigment epithelium-derived factor delivered by adeno-associated virus inhibits blood-retinal barrier breakdown in diabetic rats

Visit bosterbio.com/anti-prokineticin-1-picoband-trade-antibody-pb9881-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Prokineticin 1/PROK1 Antibody™