

## Anti-MUC3/MUC3A/MUC3B Antibody Picoband™

Catalog Number: PB9955

### About MUC3B

MUC3 consists of two genes, MUC3A and MUC3B, each encoding membrane-bound mucins possessing 2 epidermal growth factor-like domains. The MUC3 gene is mapped to chromosome 7. It was showed that synthetic peptide-mediated upregulation of MUC3 dramatically inhibited adherence of enteropathogenic E. coli or enterohemorrhage E. coli serotype O157:H7 to HT-29 human intestinal epithelial cells. Peptide stimulation altered expression of a number of transcription factors, including upregulation of SP1, CREB1, and CDX2. These transcription factors bound to consensus sites in the MUC3 promoter upon peptide stimulation and likely mediated MUC3 upregulation.

### Overview

Product Name	Anti-MUC3/MUC3A/MUC3B Antibody Picoband™
Reactive Species	Human
Description	Boster Bio Anti-MUC3/MUC3A/MUC3B Antibody Picoband™ catalog # PB9955. Tested in WB applications. This antibody reacts with Human.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q9H195

### Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human MUC3.
Predicted Reactive Species	Hamster
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.5ug/ml, Human</p>

## Anti-MUC3/MUC3A/MUC3B Antibody Picoband™ (PB9955) Images




Figure 1. Western blot analysis of MUC3 using anti-MUC3 antibody (PB9955).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human SW620 whole cell lysates,

Lane 2: human COL320 whole cell lysates,

Lane 3: human CACO-2 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes.

Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MUC3 antigen affinity purified polyclonal antibody (Catalog # PB9955) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for MUC3 at approximately 345 kDa. The expected band size for MUC3 is at 345 kDa.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-MUC3/MUC3A/MUC3B Antibody™