

Anti-Carboxypeptidase B2/CPB2 Antibody Picoband™

Catalog Number: PB9998

About Cpb2

Carboxypeptidase B2 (CPB2), also known as carboxypeptidase U (CPU), plasma carboxypeptidase B (pCPB) or thrombin-activatable fibrinolysis inhibitor (TAFI), is an enzyme that, in humans, is encoded by the gene CPB2. CPB2 is synthesized by the liver and circulates in the plasma as a plasminogen-bound zymogen. When it is activated by proteolysis at residue Arg92 by the thrombin/thrombomodulin complex, CPB2 exhibits carboxypeptidase activity. Activated CPB2 reduces fibrinolysis by removing the fibrin C-terminal residues that are important for the binding and activation of plasminogen.

Overview

Product Name	Anti-Carboxypeptidase B2/CPB2 Antibody Picoband™
Reactive Species	Mouse
Description	Boster Bio Anti-Carboxypeptidase B2/CPB2 Antibody Picoband™ catalog # PB9998. Tested in WB applications. This antibody reacts with Mouse.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q9JHH6

Technical Details

Immunogen	E. coli-derived mouse CPB2 recombinant protein (Position: K165-D387). Mouse CPB2 shares 86.1% and 96.9% amino acid (aa) sequence identity with human and rat CPB2, respectively.
Predicted Reactive Species	Bovine
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.5ug/ml, Mouse</p>

Anti-Carboxypeptidase B2/CPB2 Antibody Picoband™ (PB9998) Images



Figure 1. Western blot analysis of CPB2 using anti-CPB2 antibody (PB9998).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: mouse liver tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CPB2 antigen affinity purified polyclonal antibody (Catalog # PB9998) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for CPB2 at approximately 48 kDa. The expected band size for CPB2 is at 49 kDa.

1 Publications Citing This Product

1. PubMed ID: 19018804, INVOLVEMENT OF PROLYLCARBOXYPEPTIDASE IN THE EFFECT OF RUTAECARPINE ON THE REGRESSION OF MESENTERIC ARTERY HYPERTROPHY IN RENOVASCULAR HYPERTENSIVE RATS

Visit bosterbio.com/anti-cpb2-picoband-trade-antibody-pb9998-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Carboxypeptidase B2/CPB2 Antibody™