

Mouse recombinant IL-17D (Interleukin-17D) protein, AF

Catalog Number: PROTA0A0B4J1G4

Overview

Product Name	Mouse recombinant IL-17D (Interleukin-17D) protein, AF
Description	Interleukin 17D (IL-17D) belongs to the IL-17 family of cytokines, predicts a molecular mass of 20 kDa. IL-17 family is closely linked to host defense and immune response, and IL-17D is a novel cytokine in the IL-17 family of cytokines that has not been extensively investigated. It is highly secreted by fibrosarcoma tumor cells; in addition, ectopic expression of IL-17D in tumor cells recruits natural killer cells via the CCL2 production of endothelial cells.
Size	5ug,20ug,100ug
Tag	His Tag (N-term)
Form	Lyophilized
Source	Escherichia coli
Formulation	The protein was lyophilized from a 0.2 um filtered solution containing 20 mM sodium citrate, 0.2 M NaCl, pH 4.5. If you have any concerns or special requirements, please confirm with us.

Storage

Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

Purity

>98% as determined by SDS-PAGE.

Molecular weight

The protein has a calculated MW of 20.74 kDa. The protein migrates about 25 kDa under reducing condition (SDS-PAGE analysis).

Activity

Testing in process

Endotoxin

<0.1 EU per 1 ug of the protein by the LAL method.

Amino Acid Sequence

ALRTGRRPARPRDCADRPEELLEQLYGRLAAGVLSAFHHTLQLGPREQARNASCPAGGRAADRRFRPPTNLRVSPWAYRISYDPA RFPRYLPEAYCLCR
GCLTGLYGEEDFRFRSTPVFSPA VVLRRTAACAGGRSVYAEHYITIPVGCTCVPEPDKSAZDSANSSMDKLLLGPADRPAGR with polyhistidine tag at
the N-terminus.

Reconstitution

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 100 ug/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

Usage

Boster's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Mouse recombinant IL-17D (Interleukin-17D) protein, AF