

Human recombinant IL-7 (Interleukin-7) protein, AF

Catalog Number: PROTP13232-9

Overview

| Product Name | Human recombinant IL-7 (Interleukin-7) protein, AF |
|--------------|---|
| Description | Interleukin-7 (IL-7) is a multipotent cytokine belonged to one of the members of IL-2 superfamily. IL-7 has diverse effects on the hematopoietic and immune regulations. IL-7 is a trophic factor that is necessary for both B cell and T cell proliferation and development. In addition, it presents potential antitumor effects in tumors such as glioma, melanoma, lymphoma, leukemia, prostate cancer, and glioblastoma. |
| Size | 5ug,20ug,100ug |
| Tag | His Tag (C-term) |
| Form | Lyophilized |
| Source | Escherichia coli |
| Formulation | The protein was lyophilized from a 0.2 um filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us. |

Storage

Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

Purity

>95% as determined by SDS-PAGE.

Molecular weight

The protein has a calculated MW of 18.3 kDa. The protein migrates as 19 kDa under reducing condition (SDS-PAGE analysis).

Activity

Measured in a cell proliferation assay using PHA-activated human peripheral blood lymphocytes (PBMC). The EDII for this effect is 1 x 10 IU/mg.

Endotoxin

< 0.1 EU per 1 ug of the protein by the LAL method.







Amino Acid Sequence

MDCDIEGKDGKQYESVLMVSIDQLLDSMKEIGSNCLNNEFNFFKRHICDANKEGMFLFRAARKLRQFLKMNSTGDFDLHLLKVSEGTTILLNCTGQVK GRKPAALGEAQPTKSLEENKSLKEQKKLNDLCFLKRLLQEIKTCWNKILMGTKEH with polyhistidine tag at the C-terminus.

Reconstitution

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H[®]O to a concentration not less than 100 ug/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.



Usage

Boster's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.





Human recombinant IL-7 (Interleukin-7) protein, AF