

Mouse recombinant CCL4 (C-C Motif Chemokine Ligand 4) protein, AF

Catalog Number: PROTP14097-2

Overview

Product Name	Mouse recombinant CCL4 (C-C Motif Chemokine Ligand 4) protein, AF
Description	C-C Motif Chemokine Ligand 4 (CCL4) is a 7.66 kDa cytokine with 69 amino acid residues. CCL4, also named macrophage inflammatory protein-1beta (MIP-1beta), is mainly secreted from neutrophils, monocytes, B cells, T cells, fibroblasts, endothelial cells, and epithelial cells. In addition, CCL4 participates in immune responses, including recruitment of immune cells like lymphocytes, monocytes, and leukocytes, response to IL-1 and IFNgamma, and production of TNF when CCL4 binds to CCR5.
Size	5ug,20ug,100ug
Tag	His Tag (N-term)
Form	Lyophilized
Source	Escherichia coli
Formulation	The protein was lyophilized from a 0.2 um filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.

Storage

Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

Purity

>98% as determined by SDS-PAGE.

Molecular weight

The protein has a calculated MW of 8.64 kDa. The protein migrates as 11-17 kDa under reducing condition (SDS-PAGE analysis).

Activity

Measure by its ability to chemoattract human PBMCs using a concentration range of 20.0 - 200.0 ng/mL. Note: Results may vary from different PBMC donors.

Endotoxin

<0.1 EU per 1 ug of the protein by the LAL method.

Amino Acid Sequence

APMGSDPPTSCCFSYTSRQLHRSFVMDYYETSSLCSKPAVVFLTKRGRQICANPSEPWWTEYMSDLELN with polyhistidine tag at the N-terminus.

Reconstitution

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 100 ug/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex!

Vigorous shaking may impair the biological activity of the protein.

Usage

Boster's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Mouse recombinant CCL4 (C-C Motif Chemokine Ligand 4) protein, AF