

Human recombinant HMGB2 (High mobility group box 2) protein, AF

Catalog Number: PROTP26583-2

Overview

Product Name	Human recombinant HMGB2 (High mobility group box 2) protein, AF
Description	High mobility group protein B2, also known as HMGB2, is a member of the non-histone chromosomal high-mobility group protein family. High mobility group protein B2 is expressed as 25 kDa containing 209 amino acid residues. The function of High mobility group protein B2 is involved in transcription, chromatin remodeling and V(D)J recombination. In vitro studies have demonstrated that this protein is able to efficiently bend DNA and form DNA circles.
Size	5ug,20ug,100ug
Tag	His Tag (C-term)
Form	Lyophilized
Source	Escherichia coli
Formulation	The protein was lyophilized from a 0.2 um filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.

Storage

Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

Purity

>98% as determined by SDS-PAGE.

Molecular weight

The protein has a calculated MW of 24.84 kDa. The protein migrates as 34 kDa under reducing condition (SDS-PAGE analysis).

Activity

Testing in process

Endotoxin

< 0.1 EU per 1 ug of the protein by the LAL method.







Amino Acid Sequence

MGKGDPNKPRGKMSSYAFFVQTCREEHKKKHPDSSVNFAEFSKKCSERWKTMSAKEKSKFEDMAKSDKARYDREMKNYVPPKGDKKGKKKDPNAP KRPPSAFFLFCSEHRPKIKSEHPGLSIGDTAKKLGEMWSEQSAKDKQPYEQKAAKLKEKYEKDIAAYRAKGKSEAGKKGPGRPTGSKKKNEPEDEEEEEE EEDEDEEEEDEDEE with polyhistidine tag at the C-terminus.

Reconstitution

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H[®]O to a concentration not less than 100 ug/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.



Usage

Boster's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.





Human recombinant HMGB2 (High mobility group box 2) protein, AF