

Human recombinant 4-1BBL (4-1BB ligand) protein, AF

Catalog Number: PROTP41273-3

Overview

Product Name	Human recombinant 4-1BBL (4-1BB ligand) protein, AF
Description	4-1BB ligand (4-1BBL) is a type II transmembrane protein that is part of the tumor necrosis factor (TNF) ligand family. As an inducible co-stimulatory molecule, it presents on several antigen presenting cell (APC) types, including B cells, macrophages and DCs. The interactions between 4-1BB and 4-1BBL trigger pleiotropic effects on the immune response including antigen presenting process, proliferation of CD4 and CD8 positive T-cells, as well as cytokine secretion from T-cells through NFkappaB, c-Jun, and p38 downstream signal pathways activation. Therefore, 4-1BB and 4-1BBL are recently used for the immunotherapy of cancer.
Size	5ug,20ug,100ug
Tag	His Tag (C-term)
Form	Lyophilized
Source	Escherichia coli
Formulation	The protein was lyophilized from a 0.2 um filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.

Storage

Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

Purity

>95% as determined by SDS-PAGE.

Molecular weight

The protein has a calculated MW of 20.4 kDa. The protein migrates as 20 kDa under reducing condition (SDS-PAGE analysis).

Activity

Measure by its ability to induce IL-8 secretion in human PBMCs. The ED₅₀ for this effect is 1-5 ng/mL.

Endotoxin

<0.1 EU per 1 ug of the protein by the LAL method.

Amino Acid Sequence

MREGPELSPDDPAGLLDLRQGMFAQLVAQNVLLIDGPLSWYSDPGLAGVSLTGGLSYKEDTKELVVAKAGVYYVFFQLELRRVVAGEGSGSVSLALHLQ
PLRSAAGAAALALTVDLPPASSEARNSAFGFQGRLHLHSAGQRLGVHLHTEARARHAWQLTQGATVLGLFRVTPEIPAGLPSRSE with polyhistidine
tag at the C-terminus.

Reconstitution

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 100 ug/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

Usage

Boster's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Human recombinant 4-1BBL (4-1BB ligand) protein, AF