

Human recombinant Galectin-10 protein, AF

Catalog Number: PROTQ05315-2

Overview

Product Name	Human recombinant Galectin-10 protein, AF
Description	Galectin-10 (Gal-10) is a lectin family member and is one of the prototype galectins. It contains one carbohydrate recognition domain (CRD), responsible for beta-galactoside binding, and is biologically active as homodimers. It has been reported that galectin-10 is presented in eosinophils but also detected in basophils and macrophages. Accumulated evidence indicates that galectin-10 spontaneously forms Charcot-Leyden crystals (CLCs), participating in allergic responses and related inflammatory reactions. Several binding partners for galectin-10 have been identified, including eosinophil granule cationic ribonucleases, eosinophil-derived neurotoxin (EDN, RNS2), and eosinophil cationic protein (ECP, RNS3), demonstrating the interaction is essential for eosinophil differentiation and granulogenesis.
Size	5ug,20ug,100ug
Tag	His Tag (N-term)
Form	Lyophilized
Source	Escherichia coli
Formulation	The protein was lyophilized from a 0.2 um filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.

Storage

Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

Purity

>98% as determined by SDS-PAGE.

Molecular weight

The protein has a calculated MW of 16.31 kDa. The protein migrates as 16-19 kDa under reducing condition (SDS-PAGE analysis).

Activity

Testing in process

Endotoxin

<0.1 EU per 1 ug of the protein by the LAL method.

Amino Acid Sequence

SLLPVPTYEAASLSTGSTVTIKGRPLACFLNEPYLQVDFHTEMKEESDIVFHFQVCFGRRVVMNSREYGAWKQQVESKNMPFQDGQEFELSISVLPDKY
QVMVNGQSSYTFDHRIKPEAVKMVQVWRDISLTKFNVSYLKR with polyhistidine tag at the N-terminus.

Reconstitution

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 100 ug/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex!
Vigorous shaking may impair the biological activity of the protein.

Usage

Boster's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Human recombinant Galectin-10 protein, AF