

# Human recombinant IL-31 (Interleukin-31) protein, AF

Catalog Number: PROTQ6EBC2-2

### Overview

Product Name	Human recombinant IL-31 (Interleukin-31) protein, AF
Description	Interleukin 31 (IL-31) is an 18 kDa cytokine with 142 amino acid residues. IL-31 is mainly secreted from activated CD4+ T cells and binds to a receptor called IL-31RA. Upon binding to 31RA, IL-31 activates several signal pathways like MAPK, PI3K/AKT, and Jak/STAT pathways. It is critical in regulating many biological functions, such as cell proliferation, hematopoiesis, induction of cytokines, inflammation, and immune response.
Size	5ug,20ug,100ug
Tag	His Tag (C-term)
Form	Lyophilized
Source	Escherichia coli
Formulation	The protein was lyophilized from a 0.2 um filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.

# **Storage**

Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

# **Purity**

>98% as determined by SDS-PAGE.

### Molecular weight

The protein has a calculated MW of 16.78 kDa. The protein migrates as 18 kDa under reducing condition (SDS-PAGE analysis).

### **Activity**

Measured by its ability to induce A549 cells proliferation. The ED?? for this effect is < 40 ng/mL

### Endotoxin

< 0.1 EU per 1 ug of the protein by the LAL method.







### **Amino Acid Sequence**

MSHTLPVRLLRPSDDVQKIVEELQSLSKMLLKDVEEEKGVLVSQNYTLPCLSPDAQPPNNIHSPAIRAYLKTIRQLDNKSVIDEIIEHLDKLIFQDAPETNIS VPTDTHECKRFILTISQQFSECMDLALKSLTSGAQQATT with polyhistidine tag at the C-terminus.

### Reconstitution

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H20 to a concentration not less than 100 ug/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.



# Usage

Boster's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

# Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.





Human recombinant IL-31 (Interleukin-31) protein, AF