

Human recombinant IL-29 (Interleukin-29) protein, AF

Catalog Number: PROTQ8IU54-7

Overview

Product Name	Human recombinant IL-29 (Interleukin-29) protein, AF
Description	Interleukin 29 (IL-29) is a cytokine, predicts a molecular mass of 21.9 kDa. It belongs to type III interferons group, also termed interferons lambda (IFN-lambda). Its induction of STAT3-STAT5 has also been displayed, albeit to a lesser degree. The STAT1/STAT2 signaling cascade transpires as follows: once tyrosine residues on STAT1 and STAT2 are phosphorylated, these proteins dimerize and are subsequently transported to the nucleus.
Size	5ug,20ug,100ug
Tag	His Tag (C-term)
Form	Lyophilized
Source	Escherichia coli
Formulation	The protein was lyophilized from a 0.2 um filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.

Storage

Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

Purity

>98% as determined by SDS-PAGE.

Molecular weight

The protein has a calculated MW of 20.70 kDa. The protein migrates as 22 kDa under reducing condition (SDS-PAGE analysis).

Activity

Measure by its ability to induce IL-8 secretion in HuH7 cells. The ED₅₀ for this effect is <6 ng/mL.

Endotoxin

<0.1 EU per 1 ug of the protein by the LAL method.

Amino Acid Sequence

MPTSKPTTTGKGCHIGRFKSLSPQELASFKKARDALEESLKLKNWSCSSPVFPGNWDLRLLQVRERPVALEAELALTLKVLEAAAGPALEDVLDQPLHTLHHILSQLQACIQPQPTAGPRPRGRLHHLHRLQEAPKKESAGCLEASVTFNLFRLLTRDLKYVADGNLCLRTSTHPEST with polyhistidine tag at the C-terminus.

Reconstitution

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 100 ug/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

Usage

Boster's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Human recombinant IL-29 (Interleukin-29) protein, AF