

**Designation: HCT-8 (HRT-18)**

CLS order number: Cryovial: 300210
Vital: 330210

Origin and General Characteristics																	
Organism:	Homo sapiens (human)																
Age:	67 years of age																
Gender:	Male																
Tissue:	Colon																
Morphology:	Epithelial																
Cell type:	Colorectal adenocarcinoma, ileocecal																
Growth Properties:	Monolayer, adherent																
Description:	The HCT-8 line is identical to the HRT-18 cell line. The cells are positive for keratin by immunoperoxidase staining.																
References:	Tompkins WA et al. Cultural and antigenic properties of newly established cell strains derived from adenocarcinomas of the human colon and rectum. J. Natl. Cancer Inst. 52: 1101-1110, 1974.																
Culture Conditions and Handling																	
Culture Medium:	DMEM:Ham's F12 medium (1:1mixture) supplemented with 2 mM L-glutamine and 5% fetal calf serum (MG-40, CLS order number 820400).																
Subculturing:	Remove medium and rinse the adherent cells using PBS without calcium and magnesium (3-5 ml PBS for T25, 5-10ml for T75 cell culture flasks). Add Accutase (1-2ml per T25, 2.5ml per T75 cell culture flask), the cell sheet must be covered completely. Incubate at ambient temperature for 8-10 minutes. Carefully resuspend the cells with medium (10 ml), centrifuge for 3 min at 300xg, resuspend cells in fresh medium and dispense into new flasks which contain fresh medium.																
Split Ratio:	A ratio of 1:4 to 1:8 is recommended																
Seeding density:	$2-4 \times 10^4$ cells/cm ²																
Fluid Renewal:	Every 2 to 3 days																
Doubling time:	About 15 hrs																
Freeze Medium:	CM-1 (CLS order number: 800125, 25ml, 800150, 50ml)																
Freezing recovery:	Fast																
Sterility:	Fluorescence (DAPI) test: negative; Mycoplasma specific PCR: negative; Bacteria specific PCR: negative																
Biosafety Level:	1																
Safety precautions:	If the cryovial is planned to be stored in liquid nitrogen and to be thawed in the future, special safety precautions should be followed: Protective gloves and clothing should be used and a facemask or safety goggles must be worn when transferring frozen samples into or removing from the liquid nitrogen tank. The removal of a cryovial from liquid nitrogen may result in the explosion of the frozen vial creating flying fragments. Caputo, J.L. Biosafety procedures in cell culture. J. Tissue Cult. Methods 11:223-227, 1988. ATCC Quality Control Methods for Cell Lines, 2nd edition, 1992.																
Special Features of the Cell Line																	
Tumorigenic:	In nude mice																
Viruses:	SMRV: Negative, as confirmed by Real-Time PCR. Reverse Transcriptase negative																
DNA Profile (STR):	<table> <tr> <td>Amelogenin: X,Y</td><td>WA: 18,19</td></tr> <tr> <td>CSF1PO: 12</td><td>D3S1358: 17</td></tr> <tr> <td>D13S317: 8,11</td><td>D21S11: 29,32.2</td></tr> <tr> <td>D16S539: 12,13</td><td>D18S51: 11,17</td></tr> <tr> <td>D5S818: 13</td><td>Penta E: 7,14</td></tr> <tr> <td>D7S820: 10,12</td><td>Penta D: 9,14</td></tr> <tr> <td>TH01: 7,9.3</td><td>D8S1179: 15</td></tr> <tr> <td>TPOX: 8,11</td><td>FGA: 22</td></tr> </table>	Amelogenin: X,Y	WA: 18,19	CSF1PO: 12	D3S1358: 17	D13S317: 8,11	D21S11: 29,32.2	D16S539: 12,13	D18S51: 11,17	D5S818: 13	Penta E: 7,14	D7S820: 10,12	Penta D: 9,14	TH01: 7,9.3	D8S1179: 15	TPOX: 8,11	FGA: 22
Amelogenin: X,Y	WA: 18,19																
CSF1PO: 12	D3S1358: 17																
D13S317: 8,11	D21S11: 29,32.2																
D16S539: 12,13	D18S51: 11,17																
D5S818: 13	Penta E: 7,14																
D7S820: 10,12	Penta D: 9,14																
TH01: 7,9.3	D8S1179: 15																
TPOX: 8,11	FGA: 22																
Isoenzymes:	AK-1, 1; ES-D, 1-2; GLO-1, 2; G6PD, B; PGM1, 1; PGM3, 1; Me-2, 1																

Cell Marker:	CDx (+/-), CDy (-),
Products :	carcinoembryonic antigen (CEA) 0.5 ng/10 exp6 cells/10 days; alkaline phosphatase; keratin
Mutational profile:	HRT-18 cells carry a mutation in codon 13 of Kras gene: GGC(Wt Gly) >GAC(Asp)

Certificate of Analysis:	The Certificate of Analysis for each batch can be requested by e-mail at service@clsgmbh.de .
--------------------------	---

Recommendations for handling of adherent cell cultures following delivery

Cryopreserved cells	If immediate culturing is not intended, the cryovial(s) must be stored below -150°C or at least at -80°C after arrival. If immediate culturing is intended, please follow these instructions: Quickly thaw by rapid agitation in a 37°C water bath within 40-60 seconds. The water bath should have clean water containing an antimicrobial agent. As soon as the sample has thawed, remove the cryovial from the water bath. Note: A small ice clump should still remain and the vial should still be cold. From now on, all operations should be carried out under aseptic conditions. Transfer the cryovial to a sterile flow cabinet and wipe with 70% alcohol. Carefully open the vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of culture medium (room temperature). Resuspend the cells carefully. Centrifuge at 300xg for 3 min and discard the supernatant. The centrifugation step may be omitted, but in this case the remains of the freeze medium have to be removed 24 hours later. Resuspend the cells carefully in 10ml fresh cell culture medium and transfer them into two T25 cell culture flasks. All further steps are described in the Subculture section.
Proliferating Cultures	The cell culture flasks are completely filled with cell culture medium to prevent loss of cells during transit. Remove the entire medium except for a sufficient volume to cover the floor of the flask. Incubate at 37°C for 24 hrs. Sometimes the cultures are handled roughly during transit, and most of the cells detach and float in the culture medium. If this has occurred remove the entire content of the flask and centrifuge at 300x g for 5 minutes. Take off the supernatant, resuspend the cells in 10 ml of culture medium and transfer the entire cell suspension into cell culture flasks of suitable size (size (do not seed in more than 1T75 flask).

Warranty:	CLS warrants for a high cell viability and culture performance only if the product(s) is (are) stored and cultured according to the information described above. Using cell culture media and supplements other than the ones recommended in this product information may result in satisfactory proliferation and viabilities. CLS, however, does not warrant for cell recovery, proliferation and function if differing formulations are employed.
Disclaimer:	The customer shall not be entitled to employ this product for purposes other than research. Commercial utilization shall not be permitted; in particular, the cell line, its components or materials made therefrom shall not be sold or transferred to any third party. In addition, the term 'Commercial use' shall mean any activity by a party for consideration and may include, but is not limited to, use of the product or its components in manufacturing, for providing services, e.g. fee for service testing, in quality control or assurance processes within the manufacturing of products for sale, for therapeutic, diagnostic or prophylactic purposes, or for resale.