

NCI-H2452 Cells | 300391

General information

Description

The NCI-H2452 cell line is a human malignant pleural mesothelioma cell line, which was derived from the pleura of a patient with mesothelioma. It is frequently used in research focused on understanding mesothelioma pathophysiology and developing new therapeutic approaches. Like other mesothelioma cell lines, NCI-H2452 is associated with exposure to asbestos fibers, a well-established risk factor for mesothelioma. Studies involving NCI-H2452 have highlighted its utility in exploring mechanisms of disease progression and response to various therapies, particularly gene therapies and viral oncolysis approaches.

NCI-H2452 cells express Coxsackie and adenovirus receptor (CAR) and CD46, which makes them suitable candidates for adenovirus-based gene therapy studies. In research investigating oncolytic virotherapy, both adenovirus type 5 (Ad5) and a fiber-modified variant (Ad5F35) have been tested on NCI-H2452 cells. These adenoviruses replicate selectively within tumor cells, inducing oncolysis in a viral particle-dependent manner. It was found that both Ad5 and Ad5F35 displayed similar efficacy in inducing cell death in NCI-H2452 cells, supporting their potential in gene therapy for malignant mesothelioma.

In addition to its role in oncolytic virotherapy, NCI-H2452 cells have been used to study tumor angiogenesis, a key factor in mesothelioma progression. NCI-H2452 expresses progranulin (PGRN) and granulin-like proteins, which have been identified as novel angiogenic factors that operate independently of the VEGF pathway. This VEGF-independent angiogenesis is crucial, as it offers alternative therapeutic targets in cases where anti-VEGF therapies like bevacizumab fail to improve patient outcomes. Research indicates that these granulins contribute significantly to the formation of new blood vessels, which supports tumor growth and may be involved in the resistance to certain treatments.

Organism

Human

Tissue

Lung

Disease

Pleural biphasic mesothelioma

Synonyms

NCI-H2452, H-2452, NCIH2452

Characteristics

Age

Adult

Gender

Male

Ethnicity

European

Morphology

Epithelial

Growth properties

Adherent

NCI-H2452 Cells | 300391

Identifiers / Biosafety / Citation

Citation	NCI-H2452 (Cytion catalog number 300391)
-----------------	--

Expression / Mutation

Handling

Culture Medium	RPMI 1640, w: 2.1 mM stable Glutamine, w: 2.0 g/L NaHCO ₃ (Cytion article number 820700a)
-----------------------	--

Medium supplements	Supplement the medium with 10% FBS
---------------------------	------------------------------------

Passaging solution	Accutase
---------------------------	----------

Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
---------------------	---

Freeze medium	CM-1 (Cytion catalog number 800100)
----------------------	-------------------------------------

NCI-H2452 Cells | 300391

Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

NCI-H2452 Cells | 300391

STR profile	Amelogenin: x,y
	CSF1PO: 11,12
	D13S317: 12
	D16S539: 11,13
	D5S818: 11,12
	D7S820: 9,11
	TH01: 6,9.3
	TPOX: 8,11
	vWA: 17,18
	D3S1358: 17
	D21S11: 28,32.2
	D18S51: 15
	Penta E: 12,15
	Penta D: 9
	D8S1179: 10
	FGA: 23
	D6S1043: 11,12
	D2S1338: 20
	D12S391: 17.3,21
	D19S433: 13