

NCI-H358 Cells | 300430

General information

Description

NCI-H358, also known as H-358 or NCIH358, is an epithelial-like cell line derived from a patient with bronchioalveolar carcinoma, a subtype of non-small cell lung cancer (NSCLC). These cells display ultrastructural characteristics typical of Clara cells, such as specific cytoplasmic features. NCI-H358 cells are particularly relevant in cancer research focused on NSCLC, especially for exploring the biology and treatment of lung adenocarcinomas.

This cell line is crucial for studying the effectiveness of therapies targeting the Epidermal Growth Factor Receptor (EGFR), as mutations in EGFR are a significant focus in the treatment of NSCLC. Additionally, NCI-H358 cells are valuable for investigating the role of KRAS mutations, which are prevalent in lung cancer and known to drive oncogenic activity. The study of these mutations in NCI-H358 cells helps to elucidate the molecular pathways involved in lung cancer progression and resistance to therapies.

The NCI-H358 cell line harbors a homozygous deletion of p53, a major tumor suppressor. The H358 lung cancer cell line is also used to assess the potential of novel therapeutic approaches, such as SOS1 PROTACs, aimed at targeting specific oncogenic pathways.

In summary, the NCI-H358 cell line, derived from bronchioalveolar carcinoma, is a vital tool in NSCLC research. It is instrumental for studying EGFR-targeted therapies and KRAS mutations' role in lung cancer. Its application in cancer research extends to the development of new therapeutic strategies aimed at mitigating the effects of oncogenic mutations and improving patient outcomes in lung cancer.

Organism

Human

Tissue

Lung

Disease

Minimally invasive lung adenocarcinoma

Synonyms

NCI-H358, H-358, NCIH358

Characteristics

Age

Age unspecified

Gender

Male

Ethnicity

European

Cell type

Club cell

Growth properties

Adherent

NCI-H358 Cells | 300430

Identifiers / Biosafety / Citation

Citation	NCI-H358 (Cytion catalog number 300430)
-----------------	---

Biosafety level	1
------------------------	---

Expression / Mutation

Protein expression	UGT -, GST +, PST +, p53 -
---------------------------	----------------------------

Tumorigenic	Yes, in nude mice.
--------------------	--------------------

Mutational profile	p53 homozygously deleted
---------------------------	--------------------------

Handling

Culture Medium	RPMI 1640, w: 2.1 mM stable Glutamine, w: 2.0 g/L NaHCO ₃ (Cytion article number 820700a)
-----------------------	--

Medium supplements	Supplement the medium with 10% FBS
---------------------------	------------------------------------

Passaging solution	Accutase
---------------------------	----------

Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
---------------------	---

Freeze medium	CM-1 (Cytion catalog number 800100)
----------------------	-------------------------------------

NCI-H358 Cells | 300430

Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

Product sheet

NCI-H358 Cells | 300430

STR profile	Amelogenin: x,y
	CSF1PO: 11,12
	D13S317: 8,12
	D16S539: 12,13
	D5S818: 10,12
	D7S820: 10,11
	TH01: 6
	TPOX: 8,9
	vWA: 17
	D3S1358: 14,18
	D21S11: 28,3
	D18S51: 14
	Penta E: 18
	Penta D: 10,13
	D8S1179: 13,14
	FGA: 20,21