

PC 61 5.3 Cells | 305001

General information

Description	Rattus norvegicus (B cell); Mus musculus (myeloma), rat (B cell); mouse (myeloma)
Organism	Animal cells
Tissue	Spleen
Synonyms	PC 61-5-3, PC 61.5.3, PC615.3, PC 61, TIB-222

Characteristics

Morphology	Lymphoblast-like
Cell type	B lymphocyte
Growth properties	Suspension

Identifiers / Biosafety / Citation

Citation	PC 61 5.3 (Cytion catalog number 305001)
Biosafety level	1

Expression / Mutation

Protein expression	immunoglobulin, monoclonal antibody, against mouse interleukin-2 (interleukin 2, IL-2) receptor (CD25)
---------------------------	--

Handling

Culture Medium	DMEM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 1.5 g/L NaHCO ₃ , w: 1.0 mM Sodium pyruvate (Cytion article number 820300a)
Medium supplements	Supplement the medium with 10% FBS
Subculturing	Gently homogenize the cell suspension in the flask by pipetting up and down, then take a representative sample to determine the cell density per ml. Dilute the suspension to achieve a cell concentration of 1×10^5 cells/ml with fresh culture medium, and aliquot the adjusted suspension into new flasks for further cultivation.

PC 61 5.3 Cells | 305001

Split ratio 1:2 to 1:4

Fluid renewal 2 to 3 times per week

Freeze medium CM-1 (Cytion catalog number 800100)

Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.