

TT Cells | 305027

General information

Description TT cells continuously produce high levels of calcitonin and CEA. Immunoreactive calcitonin was found to be produced in cell culture at levels of 3900 pg/million cells and 7700 pg/million cells 24 and 72 hours respectively, after a medium change. CEA was found to accumulate to greater than 27 ng/million cells over a 72 hours period. Chromosomal analysis of the cell line and tumors induced in nude mice reveal an aneuploid human karyotype with several marker chromosomes. The initial characterization studies of the TT cell line were conducted using early passage TT cells cultivated in RPMI 1640 medium supplemented with 15% fetal bovine serum and 1mM L-glutamine. It is not known if the neuropeptides reported to be produced by this cell line when it was grown in RPMI 1640 medium are also produced by the cells when they are cultured in Ham's F-12K medium. Chromosomal analysis of the cell line and tumors induced in nude mice reveal an aneuploid human karyotype with several marker chromosomes.

Organism Human

Tissue Thyroid, medulla

Disease Hereditary thyroid gland medullary carcinoma, Multiple endocrine neoplasia type 2

Synonyms MTC-TT

Characteristics

Age 77 years

Gender Female

Ethnicity European

Morphology Epithelial

Growth properties Adherent

Identifiers / Biosafety / Citation

Citation TT (Cytion catalog number 305027)

Biosafety level 1

Expression / Mutation

TT Cells | 305027

Protein expression	Calcitonin, Carcinoembryonic Antigen(CEA)
---------------------------	---

Tumorigenic	Yes
--------------------	-----

Handling

Culture Medium	Ham's F12K Medium, w: 2.0 mM L-Glutamine, w: 2.0 mM Sodium pyruvate, w: 2.5 g/L NaHCO ₃ (Cytion article number 820608a)
-----------------------	--

Medium supplements	Supplement the medium with 10% FBS
---------------------------	------------------------------------

Passaging solution	Accutase
---------------------------	----------

Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
---------------------	---

Split ratio	1:2 to 1:4
--------------------	------------

Fluid renewal	2 to 3 times per week
----------------------	-----------------------

Freeze medium	CM-1 (Cytion catalog number 800100)
----------------------	-------------------------------------

TT Cells | 305027

Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

STR profile	Amelogenin: x,x
	CSF1PO: 10,13
	D13S317: 11
	D16S539: 12,13
	D5S818: 12,13
	D7S820: 10,12
	TH01: 6,9
	TPOX: 8,11
	vWA: 16,18
	D3S1358: 15
	D21S11: 29,32.2
	D18S51: 12
	Penta E: 7,13
	Penta D: 13,13
	D8S1179: 15,16
	FGA: 21,25
	D6S1043: 12,13
	D2S1338: 17,23
	D12S391: 15,21
	D19S433: 14,15