

## RKO Cells | 305035

## General information

## Description

RKO cells are a human colorectal carcinoma cell line widely used in research related to colon cancer. They are derived from a moderately well-differentiated adenocarcinoma of the colon and are notable for their wild-type p53 status, which is uncommon among many cancer cell lines. This feature makes RKO cells particularly valuable for studying p53 functions and the cellular mechanisms of DNA repair and apoptosis in the context of colorectal cancer.

RKO cells exhibit epithelial morphology and are characterized by their genetic stability and responsiveness to a variety of genetic and pharmacological manipulations. They are utilized in studies focusing on the molecular pathways involved in cancer progression, including cell cycle regulation, signal transduction, and metastasis. RKO cells provide insights into the role of various genes and environmental factors in colorectal cancer development and offer a platform for testing the efficacy of anti-cancer drugs.

Additionally, RKO cells are used to explore the complex interactions between cancer cells and their microenvironment, as well as the immune response to tumor cells. Their sensitivity to chemotherapeutic agents and radiation makes them suitable for use in drug discovery and development, helping to identify potential therapeutic targets and evaluate new treatment strategies for colorectal cancer.

Overall, RKO cells are a fundamental resource in colorectal cancer research, contributing significantly to our understanding of the disease's molecular biology and aiding in the development of more effective treatments.

Organism	Human
----------	-------

Tissue	Colon
--------	-------

Disease	Colon carcinoma
---------	-----------------

## Characteristics

Ethnicity	African
-----------	---------

Morphology	Epithelial
------------	------------

Growth properties	Adherent
-------------------	----------

## Identifiers / Biosafety / Citation

Citation	RKO (Cytion catalog number 305035)
----------	------------------------------------

Biosafety level	1
-----------------	---

## RKO Cells | 305035

## Expression / Mutation

<b>Receptors expressed</b>	Urokinase receptor(u-PAR)
----------------------------	---------------------------

<b>Tumorigenic</b>	Yes
--------------------	-----

## Handling

<b>Culture Medium</b>	EMEM, w: 2 mM L-Glutamine, w: 1.5 g/L NaHCO <sub>3</sub> , w: EBSS, w: 1 mM Sodium pyruvate, w: NEAA (Cytion article number 820100c)
-----------------------	--

<b>Medium supplements</b>	Supplement the medium with 10% FBS
---------------------------	------------------------------------

<b>Passaging solution</b>	Accutase
---------------------------	----------

<b>Subculturing</b>	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
---------------------	---

<b>Split ratio</b>	1:2 to 1:4
--------------------	------------

<b>Fluid renewal</b>	2 to 3 times per week
----------------------	-----------------------

<b>Freeze medium</b>	CM-1 (Cytion catalog number 800100)
----------------------	-------------------------------------

### RKO Cells | 305035

#### Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

### Quality control / Genetic profile / HLA

#### Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

**RKO Cells | 305035**

---

**STR profile**

**Amelogenin:** x,x  
**CSF1PO:** 8,1  
**D13S317:** 8,11  
**D16S539:** 12,13  
**D5S818:** 11,13  
**D7S820:** 8,1  
**TH01:** 6,1  
**TPOX:** 11  
**vWA:** 15,16,17,22  
**D3S1358:** 16,19  
**D21S11:** 27,29,30  
**D18S51:** 11,12  
**Penta E:** 11,13  
**Penta D:** 10,11  
**D8S1179:** 9,13,14  
**FGA:** 20,21,22,23  
**D1S1656:** 14,17.3  
**D6S1043:** 14,19  
**D2S1338:** 16  
**D12S391:** 15,19,20  
**D19S433:** 14