

## HEC-1-A Cells | 305077

## General information

## Description

HEC-1-A cells are a well-characterized human endometrial adenocarcinoma cell line derived from the malignant tissue of a 71-year-old Caucasian woman. This cell line, established in the mid-1970s, is extensively used in gynecological cancer research, particularly for studying endometrial carcinoma.

Morphologically, HEC-1-A cells are epithelial-like and form a monolayer of polygonal cells when cultured. They exhibit a robust and adherent growth pattern, which is typical of epithelial cells originating from solid tumors. The morphological characteristics of HEC-1-A cells make them a valuable model for studying cellular behaviors that are central to cancer progression, such as adhesion, migration, and invasion.

Genotypically, HEC-1-A cells harbor several genetic aberrations that are relevant to cancer biology, including mutations in key regulatory genes like p53 and PTEN, both of which are commonly mutated in endometrial cancer. These genetic features contribute to the cells' utility in researching the molecular underpinnings of endometrial carcinogenesis and the cellular pathways leading to tumor growth and resistance to therapy.

Research using HEC-1-A cells has significantly advanced our understanding of endometrial cancer, particularly in terms of hormonal influences, genetic mutations, and responses to chemotherapeutic agents. As a result, this cell line continues to be instrumental in developing more effective diagnostic and therapeutic strategies for endometrial carcinoma.

## Organism

Human

## Tissue

Uterus, endometrium

## Disease

Endometrial adenocarcinoma

## Synonyms

Hec-1-A, HEC-1A, HEC1-A, HEC1A, Hec1A

## Characteristics

## Age

71 years

## Gender

Female

## Ethnicity

Asian

## Morphology

Epithelial

## Growth properties

Adherent

## Identifiers / Biosafety / Citation

## HEC-1-A Cells | 305077

<b>Citation</b>	HEC-1-A (Cytion catalog number 305077)
-----------------	--

<b>Biosafety level</b>	1
------------------------	---

## Expression / Mutation

<b>Receptors expressed</b>	Receptor expression: platelet activating factor(PAF)
----------------------------	--

<b>Protein expression</b>	Oncogenes: C-Fos?
---------------------------	-------------------

<b>Antigen expression</b>	Blood Type B, Rh?
---------------------------	-------------------

<b>Tumorigenic</b>	Yes
--------------------	-----

## Handling

<b>Culture Medium</b>	McCoy's 5a, w: 3.0 g/L Glucose, w: stable Glutamine, w: 2.0 mM Sodium pyruvate, w: 2.2 g/L NaHCO <sub>3</sub> (Cytion article number 820200a)
-----------------------	---

<b>Medium supplements</b>	Supplement the medium with 10% FBS
---------------------------	------------------------------------

<b>Passaging solution</b>	Accutase
---------------------------	----------

<b>Subculturing</b>	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
---------------------	---

<b>Split ratio</b>	1:2 to 1:4
--------------------	------------

<b>Fluid renewal</b>	2 to 3 times per week
----------------------	-----------------------

<b>Freeze medium</b>	CM-1 (Cytion catalog number 800100)
----------------------	-------------------------------------

### HEC-1-A Cells | 305077

#### Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

### Quality control / Genetic profile / HLA

#### Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

**Product sheet**

**HEC-1-A Cells | 305077**

---

<b>STR profile</b>	<b>Amelogenin:</b> x,x <b>CSF1PO:</b> 10,12 <b>D13S317:</b> 11 <b>D16S539:</b> 12 <b>D5S818:</b> 11,15 <b>D7S820:</b> 9,11 <b>TH01:</b> 6,7 <b>TPOX:</b> 8,11 <b>vWA:</b> 18,19 <b>D3S1358:</b> 15 <b>D21S11:</b> 30,31 <b>D18S51:</b> 16,21 <b>Penta E:</b> 11 <b>Penta D:</b> 9,12,13 <b>D8S1179:</b> 13,14 <b>FGA:</b> 21,22 <b>D6S1043:</b> 12,18 <b>D2S1338:</b> 18,19 <b>D12S391:</b> 19 <b>D19S433:</b> 13
--------------------	--