

MS751 Cells | 305115**General information****Description**

MS751 is a tumorigenic human cervical carcinoma cell line isolated from the uterus of a female patient with epidermoid carcinoma. The cells were originally obtained from a metastatic lymph node, and they form poorly differentiated epidermoid carcinoma (grade III) when xenografted into nude mice. The tumorigenic and metastatic nature of MS751 cells makes them a valuable model for studying the processes involved in cervical cancer metastasis and tumor progression. These cells are particularly useful for investigating epithelial-to-mesenchymal transition (EMT), invasion, and metastasis, especially in relation to poorly differentiated carcinoma.

One of the key molecular features of MS751 is the presence of human papillomavirus (HPV) sequences. Originally reported to contain HPV-18, more recent studies have demonstrated that MS751 cells contain partial sequences of HPV-45, particularly from the E6/E7 region, which are expressed as poly(A)⁺ RNA. The E6 and E7 oncoproteins are well-known for their roles in disrupting the tumor suppressor functions of p53 and Rb, respectively, which promote uncontrolled cell division and contribute to oncogenesis. The presence of these viral sequences makes MS751 highly relevant for studies on HPV-associated cervical cancers, and specifically for investigating how HPV-45 contributes to the malignancy of cervical cells.

MS751 cells exhibit epithelial morphology, which is characteristic of many cervical cancer cell lines. They are widely used for research into the molecular mechanisms underlying HPV-mediated carcinogenesis, as well as for drug discovery and therapeutic screening. Given their metastatic origin and the presence of HPV sequences, MS751 provides an essential model for studying the progression of cervical cancer and testing therapeutic strategies aimed at targeting both viral and tumor-related pathways.

Organism

Human

Tissue

Cervix

Disease

Human papillomavirus-related cervical squamous cell carcinoma

Metastatic site

Lymph node

Synonyms

MS-751, MS 751

Characteristics**Age**

47 years

Gender

European

Morphology

Epithelial

Growth properties

Adherent

MS751 Cells | 305115**Identifiers / Biosafety / Citation****Citation** MS751 (Cytion catalog number 305115)**Biosafety level** 1**Expression / Mutation****Antigen expression** Blood Type AB, Rh?**Tumorigenic** Yes, in nude mice, forms poorly differentiated epidermoid carcinoma (grade?).**Viruses** HPV18, HPV45**Handling****Culture Medium** EMEM, w: 2 mM L-Glutamine, w: 1.5 g/L NaHCO₃, w: EBSS, w: 1 mM Sodium pyruvate, w: NEAA (Cytion article number 820100c)**Medium supplements** Supplement the medium with 10% FBS**Passaging solution** Accutase**Subculturing** Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.**Split ratio** 1:2 to 1:4**Fluid renewal** 2 to 3 times per week**Freeze medium** CM-1 (Cytion catalog number 800100)

MS751 Cells | 305115

Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.