

NCI-H1395 Cells | 305118

General information

Description

The NCI-H1395 cell line is derived from a non-small cell lung carcinoma (NSCLC) originating from a 55-year-old female patient with a history of significant tobacco use, quantified at 15 pack-years. Established in 1986, this cell line serves as an important biological model in lung cancer research, particularly for studying the pathophysiology and treatment responsiveness of lung adenocarcinomas. The cell line is notable for its derivation from a primary tumor, providing valuable insights into the characteristics and behavior of tumor cells directly from the patient.

Organism

Human

Tissue

Lung

Disease

Lung adenocarcinoma

Synonyms

NCI-H1395, H-1395, NCIH1395

Characteristics

Age

55 years

Gender

Female

Ethnicity

European

Morphology

Lymphoblast

Growth properties

Adherent

Identifiers / Biosafety / Citation

Citation

NCI-H1395 (Cytion catalog number 305118)

Biosafety level

1

Expression / Mutation

Handling

NCI-H1395 Cells | 305118

Culture Medium	RPMI 1640, w: 2.1 mM stable Glutamine, w: 2.0 g/L NaHCO ₃ (Cytion article number 820700a)
-----------------------	--

Medium supplements	Supplement the medium with 10% FBS
---------------------------	------------------------------------

Passaging solution	Accutase
---------------------------	----------

Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
---------------------	---

Split ratio	1: 3 to 1: 4
--------------------	--------------

Fluid renewal	2 to 3 times per week
----------------------	-----------------------

Freeze medium	CM-1 (Cytion catalog number 800100)
----------------------	-------------------------------------

NCI-H1395 Cells | 305118

Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

NCI-H1395 Cells | 305118

STR profile	Amelogenin: x,x
	CSF1PO: 12
	D13S317: 10,14
	D16S539: 11,13
	D5S818: 12
	D7S820: 8
	TH01: 6,9.3
	TPOX: 8
	vWA: 14,17
	D3S1358: 15
	D21S11: 29
	D18S51: 12,14
	Penta E: 7
	Penta D: 12
	D8S1179: 12,14
	FGA: 18,23
	D6S1043: 18,19
	D2S1338: 17,24
	D12S391: 20,22
	D19S433: 13,16