

HMy2.CIR Cells | 305126**General information****Description**

The HMy2.CIR cell line was developed through gamma irradiation and subsequent selection for the loss of HLA class I antigen expression from the HMy.2 B lymphoblastoid cell line. This parental cell line is a fast-growing mutant derived from the ARH-77 cell line. HMy2.CIR cells are particularly valuable as hosts for transfected class I major histocompatibility antigen genes, offering a versatile platform for studying antigen presentation and immune response mechanisms.

The ARH-77 cell line, from which HMy2.CIR is ultimately derived, is known to be positive for Epstein-Barr nuclear antigen (EBNA+) and Epstein-Barr viral capsid antigen (EBVCA+). Consequently, the HMy2.CIR cell line is also presumed to be EBNA positive. This cell line is characterized by its expression of small amounts of HLA Cw4, but it does not express HLA A or B locus products. This unique antigen expression profile makes HMy2.CIR cells a useful model for immunological research, particularly in the study of HLA class I-restricted antigen processing and presentation.

Organism

Human

Tissue

B-Lymphoblast

Synonyms

Hmy.2 CIR, HMy2.CIR, C1R

Characteristics**Age**

33 years

Gender

Female

Ethnicity

Caucasian

Morphology

Lymphoblast

Growth properties

Suspension

Identifiers / Biosafety / Citation**Citation**

HMy2.CIR (Cytion catalog number 305126)

Biosafety level

2

Expression / Mutation

HMy2.CIR Cells | 305126**Handling**

Culture Medium	IMDM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 25 mM HEPES, w: 1.0 mM Sodium pyruvate, w: 3.024 g/L NaHCO ₃ (Cytion article number 820800a)
-----------------------	---

Medium supplements	Supplement the medium with 10% FBS
---------------------------	------------------------------------

Subculturing	Gently homogenize the cell suspension in the flask by pipetting up and down, then take a representative sample to determine the cell density per ml. Dilute the suspension to achieve a cell concentration of 1×10^5 cells/ml with fresh culture medium, and aliquot the adjusted suspension into new flasks for further cultivation.
---------------------	--

Split ratio	1×10^5 to 1×10^6 cells/mL
--------------------	---

Fluid renewal	2 to 3 times per week
----------------------	-----------------------

Freeze medium	CM-1 (Cytion catalog number 800100)
----------------------	-------------------------------------

HMy2.CIR Cells | 305126

Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

HMy2.CIR Cells | 305126

STR profile

Amelogenin: x,x
CSF1PO: 6,1
D13S317: 11,13
D16S539: 9,13
D5S818: 10,13
D7S820: 7,12
TH01: 8
TPOX: 8
vWA: 17
D3S1358: 16
D21S11: 29,3
D18S51: 14,16
Penta E: 12
Penta D: 10
D8S1179: 15
FGA: 20,21
D6S1043: 11,19
D2S1338: 17
D12S391: 19,2
D19S433: 14,15