

HAL-01 Cells | 305140

General information

Description

The HAL-01 cell line is derived from the peripheral blood of a female adolescent with a diagnosis of acute lymphoblastic leukemia (ALL), specifically the L2 subtype. This cell line is particularly notable for containing the t(17;19)(q22;p13) chromosomal translocation, which results in the TCF3-HLF (E2A-HLF) fusion gene. This genetic feature is critical in the study of leukemia as it influences the behavior of leukemia cells, including aspects of their growth, differentiation, and response to therapies.

The presence of the TCF3-HLF fusion gene in the HAL-01 cell line makes it an invaluable resource for oncological research, particularly for studies focused on the mechanisms of leukemogenesis and the development of targeted therapies for leukemia. The fusion protein encoded by this gene is involved in the regulation of gene transcription and has been associated with poor prognosis in patients, underscoring the importance of this cell line in therapeutic development and prognostic research in acute lymphoblastic leukemia.

Organism Human

Tissue B Cell Precursor Leukemia

Synonyms HAL01, HAL-1

Characteristics

Age 17 years

Gender Female

Morphology Lymphoblast

Growth Suspension **properties**

Identifiers / Biosafety / Citation

Citation HAL-01 (Cytion catalog number 305140)

Biosafety level 1

Expression / Mutation

Handling



HAL-01 Cells | 305140

| Culture Medium | RPMI 1640, w: 2.1 mM stable Glutamine, w: 2.0 g/L NaHCO3 (Cytion article number 820700a) |
|-----------------------|--|
| Medium supplements | Supplement the medium with 10% FBS |
| Doubling time | 48 hours |
| Subculturing | Gently homogenize the cell suspension in the flask by pipetting up and down, then take a representative sample to determine the cell density per ml. Dilute the suspension to achieve a cell concentration of 1×10^5 cells/ml with fresh culture medium, and aliquot the adjusted suspension into new flasks for further cultivation. |
| Split ratio | 1: 2 to 1: 3 |
| Fluid renewal | 2 to 3 times per week |
| Freeze medium | CM-1 (Cytion catalog number 800100) |



HAL-01 Cells | 305140

Handling of cryopreserved cultures

- 1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
- 2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
- 3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
- 4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
- 5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
- 6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
- 7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
- 8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.



HAL-01 Cells | 305140

STR profile Amelogenin: x,x

CSF1PO: 11,12 **D13S317**: 11 **D16S539**: 9,11 **D5S818**: 12,13 **D7S820**: 10,12 **TH01**: 6,8 **TPOX**: 8,11 **vWA**: 16,17 **D3S1358**: 15 **D21S11**: 29,33.2 **D18S51**: 13,14 **Penta E**: 11,17 **Penta D**: 9,1 **D8S1179**: 13,15 **FGA**: 20,22 **D6S1043**: 19 **D2S1338**: 18,24 **D12S391**: 18,21 **D19S433**: 12,13.2