

B82 Cells | 305173**General information****Description**

B82 cells were derived from mouse fatty tissue, grow adherently and show a fibroblast morphology, providing researchers with a versatile model to explore a wide range of scientific questions.

B82 cells originate from NCTC clone 929, a cell line that underwent progressive selection in BUdR (bromodeoxyuridine). As a result, B82 cells demonstrate a high cloning efficiency of 70-80%, enabling researchers to generate a substantial number of genetically identical cells for their experiments. The karyotype of B82 cells is $2n = 51$.

B82 cells are a suitable experimental model for studying cellular behaviour, signalling pathways, and tissue engineering applications. With their high cloning efficiency, researchers can confidently explore a myriad of scientific questions.

Organism

Mouse

Tissue

Skin

Synonyms

B82 (Hamprecht), B-82, GM00347, GM-347, GM0347, GM347, GM00347A, GM 0347A

Characteristics**Age**

100 days

Gender

Male

Morphology

Fibroblast

Growth properties

Adherent

Identifiers / Biosafety / Citation**Citation**

B82 (Cytion catalog number 305173)

Biosafety level

1

Expression / Mutation**Handling**

B82 Cells | 305173

Culture Medium	DMEM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 1.5 g/L NaHCO ₃ , w: 1.0 mM Sodium pyruvate (Cytion article number 820300a)
-----------------------	--

Medium supplements	Supplement the medium with 10% FBS
---------------------------	------------------------------------

Passaging solution	Accutase
---------------------------	----------

Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
---------------------	---

Split ratio	1:2 to 1:5
--------------------	------------

Fluid renewal	2 to 3 times per week
----------------------	-----------------------

Freeze medium	CM-1 (Cytion catalog number 800100)
----------------------	-------------------------------------

B82 Cells | 305173

Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.