

## DMEM:Ham's F12 (1:1), w: 3.1 g/L Glucose, w: 2.5 mM L-Glutamine, w: 15 mM HEPES, w: 0.5 mM Sodium pyruvate, w: 1.2 g/L NaHCO3 | 820400a

DMEM:Ham's F12 is a widely recognized and extensively utilized basal medium in cell culture for biological research. It serves as a fundamental source of nutrients for the growth of various mammalian cell lines, particularly when supplemented with Fetal Bovine Serum (FBS).

This unique formulation combines Dulbecco's Modified Eagle Medium (DMEM) and Ham's F-12 (Ham's Nutrient Mixture F-12) in a precise 1:1 ratio. The addition of L-glutamine further enhances its composition.

DMEM, derived from Eagle's Minimal Essential Medium (EMEM), offers an increased concentration of amino acids and vitamins compared to its predecessor. In contrast, Ham's F-12 is based on Ham's F-10 medium, providing a complementary set of essential components.

To support optimal cell growth, it is common practice to supplement DMEM:Ham's F12 with FBS at a typical concentration of 5-10%. This addition is necessary as the medium lacks growth hormones, lipids, and proteins crucial for cellular development.

DMEM:Ham's F12 incorporates a pH buffer system and is often supplemented with phenol red, a pH indicator. Cultured cells in DMEM:Ham's F12, or any medium utilizing the bicarbonate buffer system, require a controlled CO<sub>2</sub> environment of 5-10% to maintain appropriate pH levels.

### Quality Control

- Sterile-filtered

### Storage and Shelf Life

- Store at +2°C to +8°C, protected from light.
- Once opened, store at 4°C and use within 6-8 weeks.

### Shipping Conditions

- Ambient temperature

### Maintenance

- Keep refrigerated at +2°C to +8°C in the dark. Avoid freezing and frequent warming to +37°C, as it reduces product quality.
- Do not heat the medium beyond 37°C or use uncontrolled heat sources such as microwave appliances.
- If only part of the medium is to be used, remove the required amount and warm it to room temperature before use.

### Composition

Category	Components	Concentration (mg/L)
Amino Acids	Glycine	18.75
	L-Alanine	4.45
	L-Arginine HCl	147.50
	L-Asparagine H <sub>2</sub> O	7.50
	L-Aspartic Acid	6.65

**DMEM:Ham's F12 (1:1), w: 3.1 g/L Glucose, w: 2.5 mM L-Glutamine, w: 15 mM HEPES, w: 0.5 mM Sodium pyruvate, w: 1.2 g/L NaHCO3 | 820400a**

L-Cysteine HCl H <sub>2</sub> O	17.56	
L-Cystine 2 HCl	31.29	
L-Glutamic Acid	7.35	
L-Glutamine	365.00	
L-Histidine HCl H <sub>2</sub> O	31.48	
L-Isoleucine	54.47	
L-Leucine	59.05	
L-Lysine HCl	91.25	
L-Methionine	17.24	
L-Phenylalanine	35.48	
L-Proline	17.25	
L-Serine	26.25	
L-Threonine	53.45	
L-Tryptophan	9.02	
L-Tyrosine Disodium Salt	48.10	
L-Valine	52.85	
Vitamins	0.0035	
	Choline Chloride	8.98
	D-Calcium Pantothenate	2.24
	Folic Acid	2.66
	myo-Inositol	12.60

**DMEM:Ham's F12 (1:1), w: 3.1 g/L Glucose, w: 2.5 mM L-Glutamine, w: 15 mM HEPES, w: 0.5 mM Sodium pyruvate, w: 1.2 g/L NaHCO3 | 820400a**

Nicotinamide	2.02
Pyridoxine HCl	0.031
Pyridoxal HCl	2.00
Riboflavin	0.219
Thiamine HCl	2.17
Vitamin B12	0.68
Inorganic Salts	
CaCl <sub>2</sub> 2 H <sub>2</sub> O	154.50
CuSO <sub>4</sub> 5 H <sub>2</sub> O	0.0013
Fe(NO <sub>3</sub> ) <sub>3</sub> 9 H <sub>2</sub> O	0.05
FeSO <sub>4</sub> 7 H <sub>2</sub> O	0.417
KCl	311.80
MgCl <sub>2</sub> 6 H <sub>2</sub> O	61.20
MgSO <sub>4</sub>	48.84
NaCl	6996.00
NaHCO <sub>3</sub>	1200.00
Na <sub>2</sub> HPO <sub>4</sub>	71.02
NaH <sub>2</sub> PO <sub>4</sub>	54.30
ZnSO <sub>4</sub> 7 H <sub>2</sub> O	0.432
Other Components	
D-Glucose	3151.00
Hypoxanthine	2.40
HEPES	3574.50

**DMEM:Ham's F12 (1:1), w: 3.1 g/L Glucose, w: 2.5 mM L-Glutamine, w: 15 mM HEPES, w: 0.5 mM Sodium pyruvate, w: 1.2 g/L NaHCO3 | 820400a**

Linoleic Acid	0.042
Lipoic Acid	0.105
Phenol Red Sodium Salt	8.63
Putrescine 2 HCl	0.081
Sodium Pyruvate	55.00
Thymidine	0.365