

HLA-DR (L243)

Туре	Size	Catalog number
Unconjugated	100µg	103201
	500µg	103203
FITC	25 tests	103214
	100 tests	103215
	200 tests	103216
PE	25 tests	103224
	100 tests	103225
	200 tests	103226
APC	25 tests	103244
	100 tests	103245
	200 tests	103246
PerCP	25 tests	103234
	100 tests	103235
	200 tests	103236
PE-Cyanine 7	25 tests	103284
	100 tests	103285
	200 tests	103286
PerCP-Cyanine 5.5	25 tests	103264
	100 tests	103265
	200 tests	103266
mFluor™ 450	25 tests	1032144
	100 tests	1032145
	200 tests	1032146
Biotin	100µg	103251

Antigen:	HLA-DR
Immunogen:	Human B lymphocytes
Host/Isotype:	Mouse, IgG2a, κ
Reactivity:	Human, Rhesus, Cynomolgus, Baboon
Purity:	>90% pure tested via polyacrylamide gel electrophoresis (PAGE)
Formulation:	PBS, pH7.2, 0.09%NaN₃ (unconjugated, Biotin)
	PBS, pH7.2, 0.09% NaN₃ and 0.2% (w/v) BSA (conjugated)
Storage:	Store at 2-8°C and protected from prolonged exposure to light. Do not freeze.
Applications:	Flow Cytometry

↔ iFluor and mFluor are trademarks of AAT Bioquest, Inc.

Application Information

Each lot of these antibodies has been pre-titrated and tested by flow cytometric analysis of human PBMCs such that $0.5\mu g$ (unconjugated, Biotin) or $5\mu l$ (conjugated) of these products are sufficient for staining 1 million cells in a $100\mu l$ staining volume or $100\mu l$ of whole blood. It is recommended to titrate antibody reactivity empirically for optimal performance. Non-human primate cross-reactivity has been validated using Caprico's PerCP-Cyanine5.5 conjugated clone L243 product.

Caprico Biotechnologies, 400 Pinnacle Way, Suite 430, Norcross, GA 30071 Phone: +1 (678) 691-2143 Technical Support Ext 102, Fax: (678) 691-2223 Email: techsupport@capricobio.com www.CapricoBio.com



TECHNICAL DATA SHEET

Antigen Information The L243 antibody reacts with human HLA-DR antigen which is expressed on B lymphocytes, monocytes, macrophages, activated T lymphocytes, activated natural killer (NK) cells, and human progenitor cells. HLA-DR is also present on thymic epithelium, B lymphocyte dependent areas of spleen and lymph node, and on B cell lymphomas.

References

- 1. Altomonte M, et al. 2004. J Cell Physiol. 200:272.
- 2. Aoudjit F, et al. 2004. Cell Res. 299:79.
- 3. Fernandez EM, et al. 2003. Hum Immunol. 64:327.
- 4. Gross U, et al. 2006. Immunobiology. 211:807.
- 5. Meguro M, et al. 2003. Cytokine. 22:107.
- 6. Stein R, et al. 2006. Blood. 108:2736.

Terms and Conditions

This product is for research use only (RUO) and not intended for diagnostic testing.