

MKK4 Recombinant Adenovirus (Constitutively Active)

CATALOG NUMBER: ADV-161

STORAGE: -80°C

QUANTITY AND CONCENTRATION: 50 µl, 1×10^{11} VP/mL in TBS containing 10% Glycerol

Background

Recombinant adenoviruses have tremendous potential in both research and therapeutic applications. There are numerous advantages in using an adenovirus to introduce genetic material into host cells. The permissive host cell range is very wide. The virus has been used to infect many mammalian cell types (both replicative and non-replicative) for high expression of the recombinant protein. Recombinant adenoviruses are especially useful for gene transfer and protein expression in cell lines that have low transfection efficiency with liposome. After entering cells, the virus remains epichromosomal (i.e. does not integrate into the host chromosome so does not activate or inactivate host genes). Recently, recombinant adenoviruses have been used to deliver RNAi into cells.

Mitogen-activated protein kinases (MAPK), including ERK1/2, p38, and JNK1/2, are important regulators of cell function. The ERK MAPKs are most frequently activated by mitogens, whereas the JNK and p38 MAPKs are strongly responsive to stress and inflammatory signals. The MAPKs are activated through multiple intracellular phosphorylation cascade events. The core unit includes MAPKKKs and MAPKKs. MKK4 and MKK7 are two closely related dual-specificity protein kinases that activate JNK. The provided recombinant adenovirus contains constitutively active form of human MKK4 sequence. The dual phosphorylation site S257/T261 in the MKK4 (E) mutant has been changed to E257/E261.

Safety Consideration

Remember that you will be working with samples containing infectious virus. Follow the recommended NIH guidelines for all materials containing BSL-2 organisms. Always wear gloves, use filtered tips and work under a biosafety hood.

Methods

The appropriate amount of viruses used for infecting cells is critical for the outcome of your experiments. If not enough virus is used, it will not give 100% of infection. If too much virus is used, it will cause cytotoxicity or other undesired effects. The amount of adenovirus cell surface receptors vary greatly among different cell types therefore the optimal concentration differs dramatically between cell types. A range of 10-200 MOI (multiplicity of infection) is used for most cell lines, but up to 1000 MOI may be used for lymphoid cell lines.

Traditionally, Infectivity particles are measured in culture by a plaque-forming unit assay (PFU) that scores the number of viral plaques as a function of dilution. In contrast to the 10-day infection of a classical plaque assay, Cell Biolabs' QuickTiter™ Adenovirus Titer Immunoassay Kit (Cat. #VPK-109) only requires 2-day infection, and there is no agar overlay step. The kit antibody against hexon protein recognizes all serotypes of adenovirus by immunocytochemistry (see Flow Chart).

Seed 293 cells in 24 or 12-well plate for 1 hr




Prepare Adenovirus Serial Dilutions
and Infect 293 cells for 48 hrs



Anti-Hexon Immunocytochemistry Staining



Count Positive Cells and Calculate Viral Titer



References

1. Bett AJ, Haddara W, Prevec L and Graham FL. (1994) *Proc Natl Acad Sci U S A.* 91:8802-6.
2. Robbins, P. D., Tahara, H., and Ghivizzani, S. C. (1998) *Trends Biotechnol.* 16, 35-40.
3. Huang, S., Stupack, D., Mathias, P., Wang, Y., and Nemerow, G. (1997) *Proc. Natl. Acad. Sci. U S A.* 94, 8156-8161.
4. Bergelson, J. M., J. A. Cunningham, G. Droguett, E. A. Kurt-Jones, A. Krithivas, J. S. Hong, M. S. Horwitz, R. L. Crowell, and R. W. Finberg. (1997) *Science* 275:1320-1323.
5. Derijard B., Raingeaud J., Barrett T., Wu I. H., Han J., Ulevitch R. J. and Davis R. J. (1995) *Science* 267:682-5.

Recent Product Citations

1. Taniguchi, C. et al. (2007). The p85 α regulatory subunit of phosphoinositide 3-kinase potentiates c-Jun N-terminal kinase-mediated insulin resistance. *Mol. Cell Biol.* 27:2830-2840.

Warranty

These products are warranted to perform as described in their labeling and in Cell Biolabs literature when used in accordance with their instructions. THERE ARE NO WARRANTIES THAT EXTEND BEYOND THIS EXPRESSED WARRANTY AND CELL BIOLABS DISCLAIMS ANY IMPLIED WARRANTY OF MERCHANTABILITY OR WARRANTY OF FITNESS FOR PARTICULAR PURPOSE. CELL BIOLABS's sole obligation and purchaser's exclusive remedy for breach of this warranty shall be, at the option of CELL BIOLABS, to repair or replace the products. In no event shall CELL BIOLABS be liable for any proximate, incidental or consequential damages in connection with the products.

This product is for RESEARCH USE ONLY; not for use in diagnostic procedures.

Contact Information

Cell Biolabs, Inc.
7758 Arjons Drive
San Diego, CA 92126
Worldwide: +1 858-271-6500
USA Toll-Free: 1-888-CBL-0505
E-mail: tech@cellbiolabs.com
www.cellbiolabs.com

©2004-2008: Cell Biolabs, Inc. - All rights reserved. No part of these works may be reproduced in any form without permissions in writing.