

## NIH3T3/GFP Cell Line

**CATALOG NUMBER:** AKR-214


**STORAGE:** Liquid nitrogen

*Note: For best results begin culture of cells immediately upon receipt. If this is not possible, store at -80°C until first culture. Store subsequent cultured cells long term in liquid nitrogen.*

**QUANTITY & CONCENTRATION:** 1 mL,  $1 \times 10^6$  cells/mL in 70% DMEM, 20% FBS, 10% DMSO

### **Background**

NIH 3T3 cells are established from a NIH Swiss mouse embryo. These cells are highly contact inhibited and are sensitive to sarcoma virus focus formation and leukaemia virus propagation. Cells have now lost their contact inhibition. The established NIH/3T3 line was subjected to more than 5 serial cycles of subcloning in order to develop a subclone with morphologic characteristics best suited for transformation assays. The "3T3" designation refers to the abbreviation of "3-day transfer, inoculum  $3 \times 10^5$  cells." The NIH3T3 cell line is one of the most commonly used fibroblast cell lines. Our NIH3T3/GFP cell line stably expresses GFP and blasticidin-resistant genes. Both GFP and blasticidin-resistant genes are introduced into parental NIH3T3 cells using lentivirus.



**Figure 1.** NIH3T3/GFP Cell Line. Left: GFP Fluorescence; Right: Phase Contrast.

### **Quality Control**

This cryovial contains at least  $1.0 \times 10^6$  NIH3T3/GFP cells as determined by morphology, trypan-blue dye exclusion, and viable cell count. The NIH3T3/GFP cells are tested free of microbial contamination.

## **Medium**

1. Culture Medium: D-MEM (high glucose), 10% fetal bovine serum (FBS), 0.1 mM MEM Non-Essential Amino Acids (NEAA), 2 mM L-glutamine, 1% Pen-Strep, (optional) 10 µg/mL Blasticidin.
2. Freeze Medium: 70% DMEM, 20% FBS, 10% DMSO.

## **Methods**

### **Establishing NIH3T3/GFP Cultures from Frozen Cells**

1. Place 10 mL of complete DMEM growth medium in a 50-mL conical tube. Thaw the frozen cryovial of cells within 1–2 minutes by gentle agitation in a 37°C water bath. Decontaminate the cryovial by wiping the surface of the vial with 70% (v/v) ethanol.
2. Transfer the thawed cell suspension to the conical tube containing 10 ml of growth medium.
3. Collect the cells by centrifugation at 1000 rpm for 5 minutes at room temperature. Remove the growth medium by aspiration.
4. Resuspend the cells in the conical tube in 15 mL of fresh growth medium by gently pipetting up and down.
5. Transfer the 15 mL of cell suspension to a T-75 tissue culture flask. Place the cells in a 37°C incubator at 5% CO<sub>2</sub>.
6. Monitor cell density daily. Cells should be passaged when the culture reaches 95% confluence.

## **Warranty**

These products are warranted to perform as described in their labeling and in Cell Biolabs literature when used in accordance with their instructions. THERE ARE NO WARRANTIES THAT EXTEND BEYOND THIS EXPRESSED WARRANTY AND CELL BIOLABS DISCLAIMS ANY IMPLIED WARRANTY OF MERCHANTABILITY OR WARRANTY OF FITNESS FOR PARTICULAR PURPOSE. CELL BIOLABS's sole obligation and purchaser's exclusive remedy for breach of this warranty shall be, at the option of CELL BIOLABS, to repair or replace the products. In no event shall CELL BIOLABS be liable for any proximate, incidental or consequential damages in connection with the products.

***This product is for RESEARCH USE ONLY; not for use in diagnostic procedures.***

## **Contact Information**

Cell Biolabs, Inc.  
7758 Arjons Drive  
San Diego, CA 92126  
Worldwide: +1 858-271-6500  
USA Toll-Free: 1-888-CBL-0505  
E-mail: [tech@cellbiolabs.com](mailto:tech@cellbiolabs.com)  
[www.cellbiolabs.com](http://www.cellbiolabs.com)

©2010-2011: Cell Biolabs, Inc. - All rights reserved. No part of these works may be reproduced in any form without permissions in writing.