


## CD28

### Mouse Anti-Human CD28 (Clone B-T3) Azide Free mAb

<b>Catalog No.</b>	CDM171	<b>Quantity:</b>	200 µg
<b>Alternate Names:</b>	Tp44		
<b>Description:</b>	The mouse monoclonal antibody recognizes human CD28, a cell-surface protein essential for T-cell activation. CD28 provides T cells with a "second signal," or co-stimulation, which, along with T cell receptor (TCR) ligation, induces proliferation and survival, and cytokine production. TCR stimulation without co-stimulation results in anergy of the T cell, CD28 interacts with CD80 (B7.1) and CD86 (B7.2), found on antigen presenting cells (APCs).		
<b>Gene ID:</b>	940		
<b>Concentration:</b>	1.0 mg/mL		
<b>Conjugate:</b>	Unconjugated		
<b>Specificity:</b>	Recognizes CD28 (the B7/BB1 receptor), a 44 kDa protein.		
<b>Host:</b>	Mouse		
<b>Isotype:</b>	IgG2a		
<b>Immunogen:</b>	U266 cell line		
<b>Clone:</b>	B-T3		
<b>Biological Activity:</b>	Inhibits mixed lymphocyte culture, co-stimulates T cell proliferation.		
<b>Formulation:</b>	Liquid in PBS, sterile filtered. Carrier and preservative free		
<b>Purification:</b>	Ion exchange chromatography		
<b>Applications:</b>	Flow Cytometry, Functional Studies, Immunohistochemistry		
<b>Application Notes:</b>	The optimal concentration should be determined by the user for each specific application.		
<b>Storage &amp; Stability:</b>	Stable at 2-8 °C for one year. For longer storage, freeze in aliquots at -20 °C to -80°C. <b>Avoid repeated freeze-thaw cycles.</b>		

Typical staining pattern of lymphocytes with the B-T3 mAb.



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

