

## PRF1

### Mouse Anti-Human Perforin Clone B-D48 Biotin Detection mAb

<b>Catalog No.</b>	CDM438	<b>Quantity:</b>	100 µg
<b>Alternate Names:</b>	P1; PFP; FLH2; PFN1; HPLH2		
<b>Description:</b>	<p>Mouse Anti-Human Perforin Clone B-D48 Biotin Detection mAb</p> <p>Background: Perforin 1 has structural and functional similarities to complement component 9 (C9). Like C9, this protein creates transmembrane tubules and is capable of lysing non-specifically a variety of target cells. This protein is one of the main cytolytic proteins of cytolytic granules, and it is known to be a key effector molecule for T-cell- and natural killer-cell-mediated cytotoxicity. Defects in this gene cause familial hemophagocytic lymphohistiocytosis type 2 (HPLH2), a rare and lethal autosomal recessive disorder of early childhood.</p>		
<b>Concentration:</b>	0.1 mg / 1.0 ml		
<b>Gene ID:</b>	5551		
<b>Specificity:</b>	Recognizes human perforin, a 70 kDa protein		
<b>Host:</b>	Mouse		
<b>Immunogen:</b>	Natural human perforin		
<b>Isotype:</b>	IgG1		
<b>Clone:</b>	B-D48		
<b>Hybridoma:</b>	Myeloma X63/AG.8653 x Balb/c spleen cells		
<b>Formulation:</b>	Liquid in PBS with 1% BSA and 0.09% Sodium Azide. Precaution: Sodium azide is a poisonous and hazardous substance which should be handled by trained staff only.		
<b>Purification:</b>	Affinity chromatography		
<b>Conjugate:</b>	Biotin		
<b>Applications:</b>	ELISA Detection Antibody. This antibody can be used as a Detection Antibody in a human Perforin sandwich Immunoassay to detect human Perforin in combination with human Perforin Capture Antibody (Cat No CDM414). The suggested coating concentration range below should be optimized by each laboratory for each application.		
<b>Application Notes:</b>	ELISA / ELISPOT: 0.05-0.5 µg/ml		
<b>Storage &amp; Stability:</b>	Store at 2-4°C for 12 months. For longer storage, freeze aliquots at -20°C. <b>Avoid repeated freeze-thaw cycles.</b>		

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

