

## BCL2

### Mouse Anti-Human BCL-2 Clone Bcl-2/100 mAb

<b>Catalog No.</b>	CMB110	<b>Quantity:</b>	100 µg
<b>Alternate Names:</b>	B-cell leukemia/lymphoma-2, B-cell lymphoma protein-2, BCL-2, PCKBCL2		
<b>Description:</b>	<p>Mouse Anti-human BCL2 Clone Bcl-2/100 monoclonal antibody. BCL2 gene was discovered by walking the chromosomal breakpoints t(14;18). This gene consists of two exons and several transcripts that encode two different protein products, BCL2α and BCL2β. Both proteins are approximately 26-30 kDa and differ at their C-terminal region. BCL2 suppresses apoptosis (programmed cell death) and this property appears to be modulated by homodimerization or association with Bax and Bad, two cell death promoters.</p>		
<b>Gene ID:</b>	596		
<b>Concentration:</b>	0.5 mg/ml		
<b>Purity:</b>	> 95%		
<b>Specificity:</b>	Recognizes the 26 kDa human BCL2		
<b>Host:</b>	Mouse		
<b>Immunogen:</b>	Synthetic peptide corresponding to residues surrounding amino acid 44 of human BCL2		
<b>Isotype:</b>	IgG1		
<b>Clone:</b>	Bcl-2/100		
<b>Formulation:</b>	Liquid in PBS + 1 mg/ml BSA + 1.5 mM sodium azide + 50% glycerol. Precaution: Sodium azide is a poisonous and hazardous substance which should be handled by trained staff only.		
<b>Purification:</b>	Protein A purified from (low FBS containing) tissue culture supernatant		
<b>Applications:</b>	<p>Western Blot                      Immunoprecipitation                      Immunohistochemistry                      Flow Cytometry</p>		
<b>Application Notes:</b>	<p>For Western Blot use a working dilution of 1-4 µg/ml.                      The optimal concentration should be determined by the user for each specific application.</p>		
<b>Storage &amp; Stability:</b>	Store at -20°C or in working aliquots at -80°C for long term storage. <b>Avoid repeated freeze-thaw cycles.</b>		

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

