

## Goat Anti-Rabbit IgG(H+L), (Ads) Biotin

<b>Catalog No.</b>	CPB015	<b>Quantity:</b>	1.0 mg
<b>Description:</b>	Goat Anti-Rabbit IgG(H+L) (Ads) Biotin. Pooled antisera from goats hyperimmunized with rabbit IgG. Cross adsorbed on mouse and human immunoglobulins and pooled sera. Purified by affinity chromatography on rabbit IgG covalently linked to agarose.		
<b>Concentration:</b>	500 µg/mL		
<b>Purity:</b>	Purified by affinity chromatography on rabbit IgG covalently linked to agarose.		
<b>Specificity:</b>	Reacts with the heavy and light chains of rabbit IgG and the light chains of rabbit IgM		
<b>Isotype:</b>	Goat IgG		
<b>Conjugate:</b>	Biotin		
<b>Formulation:</b>	Liquid volume of 2.0 mL in a buffer of PBS/NaN <sub>3</sub>		
<b>Isotype Control:</b>	Goat IgG-BIOT		
<b>Cross-Reactivity:</b>	May react with immunoglobulins from other species		
<b>Applications:</b>	ELISA, Flow Cytometry, Immunohistochemistry, Immunocytochemistry, Western Blot.		
<b>Application Notes:</b>	Working dilution for ELISA 1:5,000 - 1:20,00, Flow Cytometry $\leq 1 \mu\text{g}/10^6$ cells. Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.		
<b>Storage &amp; Stability:</b>	Store at 2-8°C. □ For long term storage at -20°C, aliquot into smaller volumes to avoid multiple freeze/thaw cycles.		

**NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.**