

## Swine Anti-Goat IgG (H+L) (Ads) Biotin pAb

<b>Catalog No.</b>	CPG006	<b>Quantity:</b>	1.0 mg
<b>Description:</b>	Swine Anti-Goat IgG (H+L) (Ads) Biotin pAb. Pooled antisera from swine hyperimmunized with goat IgG. Cross absorbed on human, rat, and mouse serum proteins. Purification by affinity chromatography on goat IgG covalently linked to agarose.		
<b>Concentration:</b>	500 µg/mL		
<b>Specificity:</b>	Reacts with the heavy and light chains of goat IgG		
<b>Immunogen:</b>	Goat IgG		
<b>Isotype:</b>	Swine IgG		
<b>Conjugate:</b>	Biotin		
<b>Formulation:</b>	Liquid volume of 2.0 mL supplied in PBS/NaN <sub>3</sub> .		
<b>Isotype Control</b>	Swine IgG-Biotin		
<b>Cross-Reactivity:</b>	May react with immunoglobulins from other species and the light chains of other goat immunoglobulins.		
<b>Applications:</b>	Applications for this product include ELISA, FLISA, Flow Cytometry, Immunohistochemistry, Immunocytochemistry, and Western Blot.		
<b>Application Notes:</b>	Working dilution for ELISA 1:5,000 - 1:20,000, for FLISA with FITC conjugate 1:200 - 1:400. Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.		
<b>Storage &amp; Stability:</b>	Store at 2-8°C. For long term storage at -20°C, aliquot into smaller volumes to avoid multiple freeze/thaw cycles.		

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

