

## Goat Anti-Human IgG (H+L) (Ads) HRP pAb

|                                 |  |                  |        |
|---------------------------------|--|------------------|--------|
| <b>Catalog No.</b>              | CPH009   | <b>Quantity:</b> | 1.0 mL |
| <b>Description:</b>             | Goat Anti-Human IgG (H+L) (Ads) HRP pAb. Pooled antisera from goats hyperimmunized with human IgG. Cross absorbed on mouse, bovine, and horse serum proteins. Purified by affinity chromatography on human IgG covalently linked to agarose. |                  |        |
| <b>Specificity:</b>             | Reacts with the heavy and light chains of human IgG and the light chains of human IgM and IgA  |                  |        |
| <b>Immunogen:</b>               | Human IgG  |                  |        |
| <b>Isotype:</b>                 | Goat IgG   |                  |        |
| <b>Conjugate:</b>               | HRP (Horseradish Peroxidase)   |                  |        |
| <b>Formulation:</b>             | Liquid volume of 1.0 mL supplied in a stock solution in 50% glycerol/50% PBS, pH 7.4. No preservative added.   |                  |        |
| <b>Isotype Control</b>          | Goat IgG-HRP   |                  |        |
| <b>Cross-Reactivity:</b>        | May react with immunoglobulins from other species.   |                  |        |
| <b>Applications:</b>            | Applications for this product include ELISA.   |                  |        |
| <b>Application Notes:</b>       | Working dilution for ELISA 1:2,000 - 1:8,000. Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.   |                  |        |
| <b>Storage &amp; Stability:</b> | Store at 2-8°C or long-term at -20°C.  |                  |        |

**NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.**

