

## Anti-Porcine Interferon-gamma (IFN- $\gamma$ ) Polyclonal Antibody

<b>Catalog No:</b>	<b>CPI009</b>
<b>Size:</b>	500 $\mu$ g
<b>Concentration:</b>	0.5 mg/0.5 mL
<b>Formulation:</b>	Rabbit polyclonal antibodies in phosphate buffered saline, pH 7.2. Carrier and preservative free. 0.22 micron sterile filtered.
<b>Endotoxin:</b>	The endotoxin level is less than 0.1 ng/ $\mu$ g of protein.
<b>Purification:</b>	Purified from rabbit serum by protein A/G affinity column chromatography.
<b>Purity:</b>	>95% as determined by SDS-PAGE analysis.
<b>Immunogen:</b>	Recombinant porcine interferon-gamma
<b>Specificity:</b>	Recombinant and natural porcine interferon-gamma
<b>Applications:</b>	This antibody has been used in Western and Dot blot applications. For Dot blot applications, a 1:10,000 dilution of the antibody will readily detect 5 ng of protein.
<b>Recommended Dilution:</b>	Centrifuge vial briefly before opening to bring contents to bottom of vial. The optimal antibody concentration should be determined for each specific application.
<b>Storage:</b>	Store at 2-8°C for up to one month. For long term use, store in aliquots at -20°C. Avoid repeated freeze/thaw cycles.
<b>Reference:</b>	1) Lee, R.S., J.R. Rusche, M.E. Maloney, D.H. Sachs, M.H. Sayegh, and J.C. Madsen (2001) CTLA4Ig-induced linked regulation of allogeneic T cell responses. <i>J. Immunol.</i> 166(3):1572-1582.

**NOT FOR HUMAN USE. FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES.**

