


## Goat Anti-Mouse IgG Fc (Ads) HRP pAb

<b>Catalog No.</b>	CPM017	<b>Quantity:</b>	1.0 mg
<b>Description:</b>	Goat Anti-Mouse IgG Fc (Ads) HRP pAb. Pooled antisera from goats hyperimmunized with mouse IgG. Cross absorbed on mouse IgG Fab, IgM, and IgA. Purified by affinity chromatography on mouse IgG covalently linked to agarose.		
<b>Concentration:</b>	500 µg/mL		
<b>Specificity:</b>	Reacts with the Fc region of mouse IgG <sub>1</sub> , IgG <sub>2a</sub> , IgG <sub>2b</sub> , IgG <sub>2c</sub> , and IgG <sub>3</sub>		
<b>Immunogen:</b>	Mouse IgG		
<b>Isotype:</b>	Goat IgG		
<b>Conjugate:</b>	HRP (Horseradish Peroxidase)		
<b>Formulation:</b>	Liquid volume of 2.0 mL supplied in a stock solution in 50% glycerol/50% PBS, pH 7.4. No preservative added.		
<b>Isotype Control</b>	Goat IgG-HRP		
<b>Cross-Reactivity:</b>	May react with immunoglobulins from other species		
<b>Applications:</b>	Applications include ELISA and Western Blot.		
<b>Application Notes:</b>	Working dilution for ELISA 1:2,000 - 1:4,000. Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.		
<b>Storage &amp; Stability:</b>	Store at 2-8°C or long-term at -20°C.		

ELISA plate was coated with purified mouse IgG Fc and IgG Fab. Immunoglobulins were detected with serially diluted Goat Anti-Mouse IgG Fc-HRP.



**NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.**