

IL3


Rat Anti-Mouse IL-3 Clone MP2-8F8 LE/NA mAb

Catalog No.	CSI10939 CSI10940	Quantity:	50 µg 0.5 mg
Alternate Names:	Interleukin-3, Burst promoting activity (BP/BPA), Eosinophil colony stimulating factor (Eo-CSF), Hematopoietic cell growth factor (HCGF), Mast cell growth factor (MGF/MCGF), Multi-colony stimulating factor (Multi-CSF), P cell stimulating activity (PCSA),		
Description:	IL-3 is a highly species-specific pleiotropic factor produced primarily by activated T cells though also by mast cells keratinocytes, and astrocytes, which stimulates colony formation of megakaryocytes, neutrophils, and macrophages from bone marrow cultures. The MP2-8F8 antibody reacts with mouse interleukin-3 (IL-3). The MP2-8F8 antibody can neutralize the bioactivity of natural or recombinant IL-3		
Concentration:	1.0 mg/ml		
Gene ID:	16187		
Host:	Rat		
Immunogen:	COS-expressed, recombinant mouse IL-3		
Isotype:	Rat IgG1, κ		
Clone:	MP2-8F8		
Bioactivity:	Proliferation/differentiation of almost all types of hematopoietic progenitors into granulocytes, macrophages, erythroid cells, megakaryocytes, mast cell colonies; induces expression of 20- α -steroid dehydrogenase, histidine, ornithine decarboxylase		
Formulation:	Phosphate-buffered solution, pH 7.2, containing no preservative. 0.2 µm filter sterilized. Endotoxin level is < 0.1 EU/µg of the protein (< 0.01 ng/µg of the protein) as determined by the LAL test.		
Purification:	The (Low Endotoxin, Azide-Free) antibody was Purified by affinity chromatography.		
Reactivity:	Mouse		
Applications:	ELISA Capture, ELISPOT Capture, ICFC, IHC, IP, Neut		
Recommended Usage:	Each lot of this antibody is quality control tested by ELISA assay. For ELISPOT applications, a concentration range of 2-8 µg/ml is recommended. For ELISA capture applications, a concentration range of 0.5-2.0 µg/ml is recommended. To obtain a linear standard curve, serial dilutions of IL-3 recombinant protein ranging from 250 to 2 pg/ml are recommended for each ELISA plate. It is recommended that the reagent be titrated for optimal performance for each application		



Storage & Stability:

The antibody solution should be stored undiluted at 4 °C. This LE/NA solution contains no preservative; handle under aseptic conditions.



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
480 Neponset Street
Bldg 12A
Canton, MA 02021

Toll Free: 888-769-1246
Phone: 781-828-0610
Fax: 781-828-0542

E-mail: info@cellsciences.com
Website: www.cellsciences.com