

IL2


Rat Anti-Mouse IL-2 Clone JES6-5H4 Biotin mAb

Catalog No.	CSI11236 CSI11237	Quantity:	50 µg 0.5 mg
Alternate Names:	Interleukin-2, T cell growth factor (TCGF), Eosinophil differentiation factor (EDF), Killer cell helper factor (KHF), Macrophage-activating factor for cytotoxicity I (MAF-C I), Thymocyte differentiation factor (TDF)		
Description:	IL-2 is a potent lymphoid cell growth factor which exerts its biological activity primarily on T cells. Additionally, IL-2 has been found to stimulate growth and differentiation of B cells, NK cells, LAK cells, monocytes, and oligodendrocytes.		
Concentration:	0.5 mg/ml		
Gene ID:	16183		
Host:	Rat		
Immunogen:	<i>E. coli</i> -expressed, recombinant mouse IL-2		
Isotype:	Rat IgG2b, κ		
Clone:	JES6-5H4		
Bioactivity:	Proliferation of T lymphocytes, B cells, anti-inflammatory, hematopoiesis, tumor surveillance		
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide. Precaution: Sodium azide is a poisonous and hazardous substance which should be handled by trained staff only.		
Purification:	The antibody was purified by affinity chromatography, and conjugated with biotin under optimal conditions. The solution is free of unconjugated biotin.		
Reactivity:	Mouse		
Applications:	ELISA Detection, ELISPOT Detection, ICFC		
Recommended Usage:	Each lot of this antibody is quality control tested by ELISA assay. For use as an ELISA detection antibody, a concentration range of 0.25-1.0 µg/ml is recommended. To obtain a		



linear standard curve, serial dilutions of IL-2 recombinant protein ranging from 250 to 2 pg/ml are recommended for each ELISA plate. For use as an ELISPOT detection antibody, a concentration range of 1-4 µg/ml is recommended. It is recommended that the reagent be titrated for optimal performance for each application

Storage & Stability: The IL-2 antibody solution should be stored undiluted at 4 °C. **Do not freeze.**



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

