

TNF


Rat Anti-Mouse TNF-alpha Clone 6B8 mAb

Catalog No.	CSI11550 CSI11551	Quantity:	50 µg 0.5 mg
Alternate Names:	Tumor necrosis factor- α , Cachectin, Necrosin, Macrophage cytotoxic factor (MCF), Differentiation inducing factor (DIF), TNFSF-2, TNF- α , TNF-alpha,		
Description:	TNF- α is secreted by macrophages, monocytes, neutrophils, T-cells (principally CD4+), and NK-cells. Many transformed cell lines also secrete TNF- α . Monomeric mouse TNF- α is 156 amino acid protein (N-glycosylated) with a reported molecular weight of 17.5 kD protein. TNF- α forms multimeric complexes; stable trimers are most common in solution. A 26 kD membrane form of TNF- α has also been described. TNF- α binding to surface receptors elicits a wide array of biologic activities including: cytolysis and cytostasis of many tumor cell lines <i>in vitro</i> , hemorrhagic necrosis of tumors <i>in vivo</i> , increased fibroblast proliferation, and enhanced chemotaxis and phagocytosis in neutrophils. The 6B8 antibody reacts with mouse and rat tumor necrosis factor- α (TNF- α).		
Concentration:	0.5 mg/ml		
Gene ID:	21926		
Host:	Rat		
Immunogen:	Recombinant mouse TNF- α		
Isotype:	Rat IgG1, κ		
Clone:	6B8		
Bioactivity:	Paracrine/endocrine mediator of inflammatory and immune functions; selectively cytotoxic for transformed cells; endothelial cell alterations; chemoattractant		
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide. Precaution: Sodium azide is a poisonous and hazardous substance which should be handled by trained staff only.		
Purification:	The antibody was purified by affinity chromatography.		
Reactivity:	Mouse		
Applications:	ELISA Capture		
Recommended Usage:	Each lot of this antibody is quality control tested by ELISA assay. For ELISA capture applications, a concentration range of 2-6 µg/ml is recommended. To obtain a linear standard curve, serial dilutions of mouse TNF- α recombinant protein ranging from 500 to 4 pg/ml are recommended for each ELISA plate. It is recommended that the reagent be titrated for optimal performance for each application.		



Storage & Stability:

The TNF- α antibody solution should be stored undiluted at 4 °C. **Avoid repeated freeze-thaw cycles**



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
480 Neponset Street
Bldg 12A
Canton, MA 02021

Toll Free: 888-769-1246
Phone: 781-828-0610
Fax: 781-828-0542

E-mail: info@cellsciences.com
Website: www.cellsciences.com