

TAF1

Rabbit Anti-Human RNAPII TBP-Associated Factor 250 kDa Clone Poly6105 pAb


Catalog No.	CSI14260 CSI14261	Quantity:	50 µl 200 µl
Alternate Names:	TAF1 RNA polymerase II, TATA box binding protein associated factor 250 kD, UBF initiation factor TFIID 250 kD subunit		
Description:	TAF250 (also known as TAF1 RNA polymerase II, TATA box binding protein associated factor 250 kD, UBF initiation factor TFIID 250 kD subunit) is a 210-250 kD member of the bipartite protein kinase family containing protein kinase domains. There are two reported isoforms of this ubiquitously expressed nuclear protein. Highest expression of TAF250 has been observed in thymus, glioblastoma, and normal endometrium at mid-secretory phase. TAF250 functions as acetyltransferase and ubiquitin activating/conjugating enzyme. This protein is thought to play an essential role in TFIID assembly by interacting with TBP and other TAFs, linking cell cycle and transcriptional control. TAF250 is essential for the progression of G1 phase of the cell cycle and has been shown to stimulate ribosomal DNA transcription by binding to the activator UBF. TAF250 is modified by phosphorylation (casein kinase II). TAF250 is the largest subunit of TFIID and has been shown to interact with TBP and other TAF proteins, and UBF. The Poly6105 antibody recognizes both isoforms of human TAF250 and has been shown to be useful for Western blotting.		
Concentration:	0.5 mg/ml		
Gene ID:	6872		
Structure:	Bipartite protein kinase family, protein kinase domains, two isoforms, 210-250 kD.		
Distribution:	Nuclear, ubiquitously expressed. Highest expression in thymus, glioblastoma, normal endometrium at mid-secretory phase.		
Host:	Rabbit		
Immunogen:	Peptide		
Isotype:	IgG		
Clone:	Poly6105		
Function:	Acetyltransferase activity; ubiquitin activating/conjugating enzyme. Thought to play an essential role in TFIID assembly by interacting with TBP and other TAFs, links cell cycle and transcriptional control.		
Formulation:	This antibody is provided in phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 50% glycerol. Precaution: Sodium azide is a poisonous and		



hazardous substance which should be handled by trained staff only.

- Purification:** The antibody was purified by antigen-affinity chromatography.
- Regulation:** Phosphorylated by casein kinase II.
- Reactivity:** Human, Recognizes both isoforms, peptide spans casein kinase II phosphorylation site
- Applications:** Western Blot
- Recommended Usage:** Each lot of this antibody is quality control tested by Western blotting. Western blotting, suggested working dilution(s): Use 10 µl per 5 ml antibody dilution buffer for each mini-gel. It is recommended that the reagent be titrated for optimal performance for each application.
- Storage & Stability:** Upon receipt, store frozen at -20° C.
- Modification:** Phosphorylation
- Interaction:** Largest subunit of TFIID, interacts with TBP and other TAF proteins, binds to UBF.

MCF7 nuclear extract was resolved by electrophoresis, transferred to nitrocellulose, and probed with rabbit anti-TAF250 antibody. Proteins were visualized using a donkey anti-rabbit secondary conjugated to HRP and a chemiluminescence detection system. In addition to the TAF250 specific band, this polyclonal antibody detects an unknown protein with a molecular weight of approximately 50 kD.



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
480 Neponset Street
Bldg 12A
Canton, MA 02021

Toll Free: 888-769-1246
Phone: 781-828-0610
Fax: 781-828-0542

E-mail: info@cellsciences.com
Website: www.cellsciences.com