

CD38, FITC-conjugated, clone HIT2 Monoclonal Antibody

Catalog No.: MON1193F

Quantity: 0.5 ml

Specificity

CD38 is present on NK-cells (LGL) and a subpopulation of T- and B-lymphocytes in peripheral blood. Furthermore it is expressed by hematopoietic precursor cells, thymocytes, and plasma cells and activated T- and B-lymphocytes. CD38 is also seen on ALL, B-cell progenitor lines and on cell lines obtained from Burkitt lymphoma.

Immunoglobulin type

Murine IgG₁

Use

The antibody can be used for flow cytometry. F/P Ratio: 3-5.

Instructions for use

This antibody has been titred for application in flow cytometry. 5 µl is the maximum amount of reagent required for 1 x 10⁶ cells. Lesser amounts of antibody may be sufficient and it is recommended that the customer determines the optimum amount of antibody for each application.

Isotype Controls

It is recommended that Monosan isotype controls be used since they have been conjugated using the same protocols as for the monoclonal antibodies: Mouse IgG₁ FITC Code M101

Presentation

100 µg / 0.5 ml antibody solution containing sodium azide. A highly purified grade of BSA has been added as a stabilizing protein to bring the final protein concentration to 4-5 mg/ml.

Minimum number of tests: 100

Storage & Handling

Store at 2-8°C for short term to 3 months and at -10°C or colder for long term storage.

Aliquot and avoid repeated freezing and thawing. Fluorochrome conjugates should be stored in the dark and direct exposure to light should be avoided during handling and cell staining.

FOR RESEARCH USE ONLY, NOT FOR DRUG, DIAGNOSTIC OR OTHER USE.



Cell Sciences, Inc.
480 Neponset Street
Bldg 12A
Canton, MA 02021

Toll Free: 888-769-1246
Phone: 781-828-0610
Fax: 781-828-0542

E-mail: info@cellsciences.com
Web Site: www.cellsciences.com