

Endothelium, clone PAL-E Monoclonal Antibody

Catalog No.: MON 6001-1

Quantity: 1 ml

Specificity

The antibody only stains human and various animal blood vessels (goat, rabbit, pig, dog) with the exception of arteries. It does not stain rat, mouse and chicken endothelium.

Reactivity with:

Lymphatic endothelium: not detectable.
Sinus histiocytes: not detectable.
Megakaryocytes: not detectable.

Immunoglobulin type

Murine IgG2_a

Use

The antibody is useful for immunohistochemistry on aceton fixed frozen sections. Not useful on cell suspensions or Western blotting.

Instructions for use

The working dilution for frozen sections is approximately 1:100, preferably in PBS.

Presentation

1 ml undiluted supernatant with approximately 15 µg antibody/ml and 10 mM sodium azide.

Literature

- Jones, R.R., et al., 1986, J. Clin. Path. 39, 742-749.
- Holden, C.A., et al., 1986, J. Immun. Meth. 91, 45-52.
- Schlingemann, R.O., et al., 1987, Laboratory Investigation 52, 71.

For research use only. Not for human use. Not for diagnostic or therapeutic use.

Toll Free: 888-769-1246

Phone: 781-828-0610

Fax: 781-828-0542

E-mail: info@cellsciences.com

Web Site: www.cellsciences.com