

Human Eosinophil Major Basic Protein, Clone BMK-13 Monoclonal Antibody

Catalog No: MON6008-1

Quantity: 1 ml

Specificity:

BMK-13 binds to the 10 kD eosinophil Major Basic Protein (MBP) of both resting and activated eosinophils in cytopins and frozen sections of bronchial and skin biopsies of allergic sites and normal sites and thus can be used as a "pan-eosinophil" marker. The antibody BMK-13 stains in frozen sections of bronchial biopsies from atopic asthmatics, rhinitics and normal non-atopic subjects, substantially higher counts of positive cells when compared to EG1, EG2 and chromotrope 2R. BMK-13 cross-reacts weakly with human basophils, which also contain low level of this protein. It does not cross-react with any other human protein or cell.

Immunoglobulin type:

Murine IgG₁

Use:

This mAb is a very useful tool for clinical and experimental allergy studies. BMK-13 is a pan-eosinophil marker in both frozen sections and paraffin embedded tissues. It stains in bronchial sections 98% of the periphery blood eosinophils. Further, the antibody can be used in specific ELISA tests to determine the release of MBP and to quantify the number of eosinophils in various *in vitro* adhesion assays.

Instructions for use:

Most useful for immunohistochemistry on frozen sections is the APAAP method. BMK-13 also binds to eosinophils in paraffin-embedded tissue. A 1:30 dilution in PBS is recommended. Incubation time: 1 hour at RT.

Protease pretreatment is necessary. We recommend using pepsine (Roche, Cat. No.108057)

DO NOT use heat denaturation. This antibody is suitable for blood smears (cytopins) if fixed with methanol-acetone mixture (1:1, v/v). It is unlikely, without permeabilization, that the BMK13 is suitable for the flow cytometry because BMK13 stains intracellular, stored proteins.

Presentation:

1 ml supernatant solution in RPMI-1640 with approximately 10 µg antibody/ml and 0.1% sodium azide. Sufficient for 50-250 tests. Catalog No. MON6008-1.

5 ml supernatant solution in RPMI-1640 with approximately 10 µg antibody/ml and 0.1% sodium azide. Sufficient for 250-1250 tests. Catalog No. MON6008-5.

Literature :

- Barkans, J.R., et al., 1990, Clin. & Exp. Allergy 20, suppl. 1, page 1.
- Moqbel, R., et al., 1992, Clin & Exp. Allergy 22, 265-273.
- Spry, C.J.F., et al., 1992, "Trends, eosinophils 1992" Immunology today 13, 384-387.

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences, Inc.
Neponset Valley Tech Park
480 Neponset St., Bldg. 12
Canton, MA 02021

Toll Free: 888 769-1246
Phone: 781 828-0610
Fax: 781 828-0542

E-mail: info@cellsciences.com
Web Site: www.cellsciences.com