

Epidermal Growth Factor, EGF, murine, clone F5 Monoclonal Antibody

Catalog No.: MON 8002-1

Quantity: 1 ml

Specificity

The antibody reacts with mouse EGF in ELISA and in spot blots. In immunohistochemistry the antibody reacts with formalin fixed and paraffin embedded mouse salivary glands. It also reacts with human Brunner's glands (presumably with urogastron).

Immunoglobulin type

Murine IgM

Use

Detection of murine EGF. In ELISA 10ng is detectable. In Western blot 1ng is detectable. Antibody can be used on formalin fixed frozen sections and paraffin sections of salivary glands.

Instructions for use

For use in staining of frozen tissue sections the working dilution is 1:20 preferably in phosphate buffered saline. See for further instructions G and H.

Presentation

1 ml tissue culture supernatant with approximately 10 μ g antibody/ml and 1% BSA and 20 mM sodium azide. Catalog No.: MON 8002-1.

5 ml tissue culture supernatant with approximately 10 μ g antibody/ml and 1% BSA and 20 mM sodium azide. Catalog No.: MON 8002-5.

Toll Free: 888-769-1246

Phone: 781-828-0610

Fax: 781-828-0542

E-mail: info@cellsciences.com

Web Site: www.cellsciences.com

FOR RESEARCH USE ONLY, NOT FOR DRUG, DIAGNOSTIC OR OTHER USE.